



# Foraminifera from the Kuromatsunai and Setana Formations in southwestern Hokkaido, Japan -Preliminary report-

メタデータ	言語: eng 出版者: 公開日: 2012-10-29 キーワード (Ja): キーワード (En): 作成者: 能條, 歩 メールアドレス: 所属:
URL	<a href="https://hokkyodai.repo.nii.ac.jp/records/9395">https://hokkyodai.repo.nii.ac.jp/records/9395</a>

# Foraminifera from the Kuromatsunai and Setana Formations in southwestern Hokkaido, Japan

## -Preliminary report-

Ayumu NOJO<sup>1)</sup>

**Key words :** foraminifera, Kuromatsunai Formation, Setana Formation, Pliocene, Pleistocene,  
southwestern Hokkaido.

### 1. Introduction

Neogene to Quaternary deposits distributed along the Japan Sea side in Hokkaido, implying a history of development and changes of the Japan Sea, are one of most important subjects for the study to understand geological and geographical frameworks of the present Japanese Islands. In spite of these facts, there have not been enough data about the environmental changes of these areas, near the northern end of the Japan Sea areas, or around the Oshima Peninsula.

The Oshima Peninsula in southwestern Hokkaido is regarded geologically as the northern extension of the "Green Tuff" region of the Inner Zone of the Northeast Japan (e.g. Kato et al. eds., 1990). The peninsula has extensive distributions of the Neogene and Quaternary deposits which unconformably overlie pre-Tertiary sedimentary and igneous rocks. This region has been one of stratigraphical standards of the Neogene in Hokkaido since the study by Nagao and Sasa (1933).

Neogene to Pleistocene marine sediments in this region are generally classified into the following ascending-ordered formations, Kunnu, Yakumo, Kuromatsunai, and Setana (Nagao and Sasa, 1933). These formations are well exposed in middle to northern parts of the peninsula. Such stratigraphy first proposed by Nagao and Sasa (1933) has been accepted, but recent biostratigraphical data reveal that the Setana Formation should be of Pleistocene age (e.g. Nojo et

<sup>1)</sup> Department of Science Education, Iwamizawa College, Hokkaido University of Education, Iwamizawa. 068-8642 Japan.

al., in press). General relationships among these formations are as follows: from the Kunnui to Kuromatsunai Formation they contact conformably each other, but they are clinounconformably overlain by the Setana (e.g. Nagao and Sasa, 1933; Matsui et al., 1955; Hashimoto et al., 1963; Oka and Mitani, 1981; Nojo et al., 1994).

There are a few reports on microfossils in the Kuromatsunai and Setana Formations. In particular, despite abundant yieldings of foraminifera from the Setana Formation, there have not been so many reports on foraminifera (Asano, 1936a, b, 1937, 1939; Asano and Nakamura, 1937; Shirai, 1959, 1960; Masatani and Ohkura, 1980MS; Takahashi et al., 1980MS; Tsuchi and Ibaragi, 1982; Tsubakihara et al., 1989; Nojo et al., 1997), and we have only chronostratigraphical data by Nojo et al.(in press), which are related to paleoenvironmental changes on the basis of micropaleontological analyses.

The rock materials for this study were sampled from the formations in northern to middle part of the Oshima Peninsula, and were analyzed in 1994 to 1998 by the author when he was a curator of Imakane Borad of Education. Some data of these works have been reported by Nojo et al. (1997: Tab. 1). This study is preliminaly report and list of foraminifera from the Kuromatsunai and Setana Formations to indicate a history of paleoenvironmental changes. The author mainly describes about some occurrences of foraminifera (Tab. 2) and shows their fuanal list with plates in this report. A detailed discussion about paleoenvironmental changes is left for another opportunity.

Table 1 Foraminifera from the Kuromatsunai and Setana Formation in Imakane area (after Nojo et al., 1997)

Sample number (Kmu: Sakurubetsu Pyro. Mem., Kms: Okusawa Sand. Mem, Ss: Setana F.)	Kmu-1	Kmu-2	Kmu-3	Kmu-4	Kmu-5	Kms-1	Ss-1	Ss-2	Sample number (Kmu: Sakurubetsu Pyro. Mem., Kms: Okusawa Sand. Mem, Ss: Setana F.)	Kmu-1	Kmu-2	Kmu-3	Kmu-4	Kmu-5	Kms-1	Ss-1	Ss-2
<b>Planktonic Foraminifera</b>																	
<i>Globigerina bulloides</i> (Orbigny)	+	+	+	+			+		<i>Fissurina cf. rizae</i> Seguenza					+			
<i>Globigerina quinqueloba</i> Naitland			+						<i>Fissurina cf. subquadra</i> Parr		+						
<i>Globigerina</i> sp.							+		<i>Fissurina</i> sp.								
<i>Globigerinella glutinata</i> (Egger)									<i>Globigerinella</i> spp.								
<i>Globigerinella uvula</i> (Ehrenberg)		+					+		<i>Globocassidulina bisecta</i> Nomura		+	+	+				+
<i>Neogloboquadrina incompta</i> (Cifelli)	+	+	+	+			+		<i>Globocassidulina jamesoni</i> (McCulloch)								
<i>Neogloboquadrina pachyderma</i> (Ehrenberg) (dextral)	+	+	+	+			+		<i>Globocassidulina</i> spp.		+						+
<i>Neogloboquadrina pachyderma</i> (Ehrenberg) (sinistral)	+	+	+	+			+		<i>Guttulina yabei</i> Cushman & Ozawa								
<b>Benthic Foraminifera</b>																	
<i>Ammonia</i> sp.									<i>Guttulina</i> spp.								
<i>Angulogerina kokuryuensis</i> Asano			+	+					<i>Hanawata nipponica</i> Asano								
<i>Angulogerina</i> sp.		+							<i>Heterolepa praecincta</i> (Karrer)								
<i>Asteronion hamadaense</i> Asano			+	+					<i>Heterolepa subhadingeri</i> (Parr)								
<i>Bolivina decussata</i> Brady			+	+					<i>Islandella japonica</i> (Asano & Nakamura)		+						
<i>Bolivina</i> spp.		+	+						<i>Islandella norcrossi</i> (Cushman)								
<i>Bolivina quadrilatera</i> (Schwager)	+								<i>Islandella setanaensis</i> (Asano & Nakamura)		+						
<i>Buccella frigida</i> (Cushman)			+	+					<i>Islandella subimbrata</i> (Asano & Nakamura)		+	+					
<i>Buccella tenurima</i> (Bandy)		+							<i>Islandella yabei</i> (Asano & Nakamura)		+	+					
<i>Buccella</i> sp.			+						<i>Lagenaria australis</i> Reuss								
<i>Buliminella</i> sp.									<i>Lagenaria apiculata</i> Loeblich & Tappan								
<i>Cibicides lobatulus</i> (Walker & Jacob)				+					<i>Lagenaria striata</i> (Orbigny)								
<i>Cibicides refidens</i> de Montfort			+						<i>Lagenaria</i> spp.								
<i>Cibicides subdepressus</i> (Asano)				+					<i>Lenticulina</i> sp.								
<i>Cibicides</i> spp.				+					<i>Melonis pomphiloides</i> (Fichtel & Moll)								
<i>Cribroelphidium yabei</i> (Asano)					+				<i>Melonis</i> sp.								
<i>Cribroelphidium</i> sp.									<i>Nonion</i> sp.								
<i>Dyocibicides perforatus</i> Cushman & Valentine	+								<i>Nonionillina labradorica</i> (Dawson)								
<i>Elphidium crispum</i> (Linné)			+						<i>Olina striatopunctata</i> (Parker & Jones)								
<i>Elphidium excavatum</i> (Terquem) forma <i>clavata</i> (Cushman)				+					<i>Oridorasalis umbonatus</i> (Reuss)								
<i>Elphidium jensenii</i> (Cushman)					+				<i>Planoglabratella subopercularis</i> (Asano)								
<i>Elphidium subarcticum</i> Cushman					+				<i>Polystomellina discorbicularis</i> Yabe & Hanzawa								
<i>Elphidium</i> spp.					+				<i>Pseudoparrella takayanagi</i> (Iwasaki)								
<i>Epistominella pulchella</i> Husezima & Maruhasi					+				<i>Pullenia apertura</i> Cushman								
<i>Epistominella</i> spp.					+				<i>Pyrgo ezo</i> Asano								
<i>Eponides</i> ? spp.					+				<i>Siliostomella</i> sp.								
<i>Fissurina amnicola</i> (Burrows & Holland)									<i>Uvigerina akitaensis</i> Asano								
<i>Fissurina baccata</i> (Hellier-Allen & Earland)	+				+				<i>Uvigerina juncea</i> Cushman & Todd								
<i>Fissurina marginata</i> (Montagu)					+				<i>Uvigerina schenki</i> Asano								
<i>Fissurina orbignyi</i> Seguenza					+				<i>Uvigerina</i> spp.								

Foraminifera from the Kuromatsunai and Setana Formations

Table 2 Foraminifera from the Kuromatsunai and Setana Formations

Sampling horizon	Upper Kuroma tsu-nal F.	Lower Setana F.	Upper Setana F.	Sampling horizon	Upper Kuroma tsu-nal F.	Lower Setana F.	Upper Setana F.
<b>Planktonic Foraminifera</b>							
<i>Globigerina bulloides</i> d'Orbigny	A	A	A	<i>Discorbella bertheloti</i> (d'Orbigny)	-	-	R
<i>Globigerina quinqueloba</i> Natland	R	C	F	<i>Discorbella</i> spp.	R	-	VR
<i>Globigerinella glutinata</i> (Egger)	F	F	F	<i>Eilohedra</i> spp.	R	-	-
<i>Globigerinella uvula</i> (Ehrenberg)	R	VR	F	<i>Elphidiella arctica</i> (Parker & Jones)	C	-	-
<i>Globigerinoides ruber</i> (d'Orbigny)	-	-	R	<i>Elphidium aculeatum</i> (d'Orbigny)	-	-	VR
<i>Globorotalia inflata</i> (d'Orbigny)	C	C	VR	<i>Elphidium advena</i> (Cushman)	-	-	VR
<i>Grobigerina</i> spp.	-	-	R	<i>Elphidium bartletti</i> Cushman	R	F	R
<i>Neogloboquadrina</i> cf. <i>asanoi</i> (Maiya, Saito, and Sato)	VR	-	-	<i>Elphidium crispum</i> (Linné)	F	F	F
<i>Neogloboquadrina dutertrei</i> (d'Orbigny)	VR	-	-	<i>Elphidium excavatum</i> (Terquem) forma <i>clavata</i> Cushman	-	R	F
<i>Neogloboquadrina incompta</i> (Cifelli)	C	A	A	<i>Elphidium frigidum</i> Cushman	R	-	R
<i>Neogloboquadrina pachyderma</i> (D) (Ehrenberg)	F	A	C	<i>Elphidium hanzawai</i> Asano	-	VR	-
<i>Neogloboquadrina pachyderma</i> (S) (Ehrenberg)	F	C	C	<i>Elphidium incertum</i> (Williamson)	-	-	VR
<i>Neogloboquadrina</i> sp.	-	VR	R	<i>Elphidium janseni</i> (Cushman)	VR	-	R
Gen. et sp. indet.	VR	R	VR	<i>Elphidium macellum</i> (Fichtel & Moll)	VR	-	-
<b>TOTAL COUNTED</b>	296	308	5035	<i>Elphidium subarcticum</i> Cushman	-	F	C
<b>Benthic Foraminifera</b>							
<i>Alabamina japonica</i> (Asano)	VR	-	-	<i>Elphidium subgranulosum</i> Asano	-	-	F
<i>Ammonia beccarii</i> (Linné)	-	VR	-	<i>Elphidium subincertum</i> Asano	-	R	-
<i>Ammonia ketienensis</i> (Ishizaki)	-	VR	-	<i>Elphidium translucens</i> Natland	-	-	R
<i>Ammonia inflata</i> (Seguenza)	-	R	-	<i>Elphidium</i> spp.	F	VR	R
<i>Ammonia takanabensis</i> (Ishizaki)	-	VR	-	<i>Epistominella pulchella</i> Husezima & Maruhashi	R	VR	R
<i>Ammonia</i> spp.	-	VR	R	<i>Epistominella</i> spp.	VR	-	R
<i>Angulogerina hugheei</i> (Galloway and Wissler)	R	VR	R	<i>Eponides?</i> sp.	R	-	R
<i>Angulogerina kokozuraensis</i> Asano	R	-	R	<i>Fissurina annectens</i> (Burrows & Holland)	-	-	R
<i>Anomalina rostata</i> (Brady)	-	-	VR	<i>Fissurina</i> cf. <i>annectens</i> (Burrows & Holland)	VR	-	-
<i>Anomalina</i> spp.	-	-	R	<i>Fissurina lacunata</i> (Burrows & Holland)	-	-	R
<i>Anomalinoides globulosus</i> (Chapman & Parr)	VR	-	R	<i>Fissurina lucida</i> (Williamson)	-	-	R
<i>Astacolus hyalaculus</i> Loeblich and Tappan	VR	-	VR	<i>Fissurina marginata</i> (Walker & Boys)	R	R	F
<i>Astrononion aomoriense</i> Asano	-	VR	-	<i>Fissurina obscurcostata</i> Galloway & Wissler	VR	-	R
<i>Astrononion hamadaense</i> Asano	-	F	R	<i>Fissurina orbigniana</i> Seguenza	R	-	VR
<i>Astrononion</i> spp.	R	-	VR	<i>Fissurina</i> cf. <i>rizzae</i> Seguenza	-	-	R
<i>Bolivina decussata</i> Brady	F	-	R	<i>Fissurina semimarginata</i> (Reuss)	VR	R	-
<i>Bolivina</i> sp. A	VR	R	R	<i>Fissurina</i> cf. <i>subquadrata</i> Parr	R	VR	R
<i>Bolivina</i> spp.	R	-	R	<i>Fissurina</i> spp.	R	VR	R
<i>Brizalina alata</i> (Seguenza)	-	-	VR	<i>Gavelinopsis praegeri</i> (Heron-Allen & Earland)	-	-	R
<i>Brizalina</i> spp.	-	-	R	<i>Gavelinopsis</i> spp.	R	-	R
<i>Buccella frigida</i> (Cushman)	VR	R	R	<i>Glabratella</i> cf. <i>aurantista</i> Seiglie & Bermúdez	-	-	R
<i>Buccella makiyamai</i> (Chihi)	VR	R	C	<i>Glabratella murabilis panamensis</i> Seiglie & Bermúdez	-	-	R
<i>Buccella nipponica</i> (Husezima & Maruhashi)	-	VR	R	<i>Glabratella pulvinata</i> (Brady)	-	-	R
<i>Buccella tanaii</i> (Uchio)	-	-	R	<i>Glabratella</i> sp. A	-	-	R
<i>Buccella tenerima</i> (Bandy)	R	-	C	<i>Glabratella</i> sp. B	-	-	R
<i>Buccella</i> spp.	VR	VR	R	<i>Glabratella</i> spp.	VR	-	R
<i>Buliminella</i> cf. <i>imamurae</i> Tai	R	-	-	<i>Glandulina ovula</i> d'Orbigny	-	-	R
<i>Buliminella elegantissima</i> (d'Orbigny)	-	-	F	<i>Globocassidulina bisecta</i> Nomura	F	R	R
<i>Cancris auriculus</i> (Fichtel & Moll)	-	-	VR	<i>Globocassidulina canalisuturata</i> Eade	-	R	-
<i>Chrysalidinella dimorpha</i> (Brady)	-	-	VR	<i>Globocassidulina neobrocha</i>	-	-	R
<i>Cibicides</i> cf. <i>kamadai</i> Asano	-	-	VR	<i>Globocassidulina</i> spp.	F	R	F
<i>Cibicides lobatulus</i> (Walker & Jacob)	A	A	F	<i>Guttulina</i> spp.	VR	-	R
<i>Cibicides refulgens</i> de Montfort	C	C	C	<i>Gyroidina</i> spp.	-	-	R
<i>Cibicides subdepressus</i> (Asano)	R	F	R	<i>Hanzawaia nipponica</i> Asano	R	VR	R
<i>Cibicides tani</i> Iwasa and Kikuchi	F	R	R	<i>Haynesina</i> sp.	-	-	R
<i>Cibicides</i> spp.	R	R	R	<i>Heronallenia</i> sp.	-	-	VR
<i>Cibicidoides pachydermus</i> (Rzehak)	R	-	VR	<i>Heterolepa subhaidingerii</i> (Parr)	-	-	R
<i>Cornuspira involvens</i> (Reuss)	-	-	R	<i>Heterolepa</i> sp.	VR	-	-
<i>Cribroelphidium oregonense</i> (Cushman & Grant)	-	-	VR	<i>Islandiella helenae</i> Feyling-Hassen & Buzas	-	-	R
<i>Cribroelphidium yabei</i> (Asano)	-	-	R	<i>Islandiella islandica</i> (Nørberg)	F	-	-
<i>Cribroelphidium curta</i> (Cushman)	-	-	VR	<i>Islandiella japonica</i> (Asano & Nakamura)	F	C	R
<i>Cycloforia contorta</i> (d'Orbigny)	-	VR	R	<i>Islandiella norcrossi</i> (Cushman)	-	-	R
<i>Cystammina pauciloculata</i> (Brady)	VR	-	-	<i>Islandiella sublimbata</i> (Asano & Nakamura)	A	A	F
<i>Discorbella araucana</i> (d'Orbigny)	R	VR	F	<i>Islandiella yabei</i> (Asano & Nakamura)	F	F	-
				<i>Islandiella</i> spp.	-	R	R

Table 2 Foraminifera from the Kuromatsunai and Setana Foramtions (continued)

Sampling horizon	Upper Kuroma tsu-nai F.	Lower Setana F.	Upper Setana F.	Sampling horizon	Upper Kuroma tsu-nai F.	Lower Setana F.	Upper Setana F.
<b>Benthic Foraminifera</b>				<b>Benthic Foraminifera</b>			
<i>Karreriella baccata</i> japonica Asano	-	VR	-	<i>Quinqueloculina akmeriana</i> d'Orbigny	R	R	R
<i>Karreriella</i> spp.	R	-	R	<i>Quinqueloculina costata</i> d'Orbigny	-	-	R
<i>Lagena acuticosta</i> (Reuss)	-	-	VR	<i>Quinqueloculina elongata</i> Natland	-	-	R
<i>Lagena apiopleura</i> (Loeblich & Tappan)	VR	VR	R	<i>Quinqueloculina kuromatnaiensis</i> Asano	-	-	R
<i>Lagena sulcata spicata</i> Cushman & McCulloch	-	-	R	<i>Quinqueloculina sawanensis</i> Asano	-	-	R
<i>Lagena</i> sp.	-	-	VR	<i>Quinqueloculina seminulum</i> (Linné)	-	VR	R
<i>Lenticulina nikobarensis</i> (Schwager)	-	-	VR	<i>Quinqueroculina vulgaris</i> d'Orbigny	-	R	R
<i>Lenticulina</i> sp.	-	-	VR	<i>Quinqueloculina</i> cf. <i>vulgaris</i> d'Orbigny	-	-	VR
<i>Melonis pacificus</i> (Cushman)	VR	-	VR	<i>Quinqueloculina yezoensis</i> Asano	-	-	R
<i>Melonis uchiori</i> Hasegawa	-	-	VR	<i>Quinqueloculina</i> sp. B	-	-	R
<i>Melonis</i> spp.	VR	-	-	<i>Quinqueloculina</i> sp. C	-	-	R
<i>Miliolinella circularis</i> (Bornemann)	-	-	R	<i>Quinqueloculina</i> sp. F	-	-	R
<i>Miliolinella</i> spp.	-	-	R	<i>Quinqueloculina</i> sp. G	-	-	R
<i>Neococonorbina stachi</i> (Asano)	-	VR	-	<i>Quinqueloculina</i> sp. H	-	-	R
<i>Neoeponides procerus</i> (Brady)	-	R	R	<i>Quinqueloculina</i> spp.	R	R	F
<i>Neoeponides</i> spp.	VR	-	-	<i>Rectobolivina rephanus</i> (Parker & Jones)	-	-	VR
<i>Nodosarina?</i> sp.	-	-	VR	<i>Rosalina australis</i> (Parr)	VR	VR	R
<i>Nomion japonicus</i> Asano	-	-	R	<i>Rosalina bradyi</i> (Cushman)	VR	VR	F
<i>Nomion</i> spp.	R	VR	R	<i>Rosalina isabelleana</i> d'Orbigny	-	-	R
<i>Nomionella stella</i> Cushman & Moyer	-	-	R	<i>Rosalina vilardeboana</i> d'Orbigny	-	-	R
<i>Nomionoides grateloupi</i> (d'Orbigny)	-	-	VR	<i>Rosalina</i> spp.	R	VR	R
<i>Colina carteri</i> Albani and Yassini	VR	-	-	<i>Sigmohauerina</i> sp.	-	-	VR
<i>Colina costata</i> (Williamson)	-	-	VR	<i>Sigmoidella pacifica</i> Cushman & Ozawa	-	-	VR
<i>Colina melo</i> d'Orbigny	-	VR	R	<i>Sigmoidella</i> spp.	VR	-	-
<i>Colina striopunctata</i> (Parker & Jones)	-	-	VR	<i>Sigmoilina sigmaidea compressa</i> Cushman	-	R	-
<i>Colina</i> sp.	-	-	VR	<i>Sigmoilina</i> sp.	-	-	VR
<i>Oridosalis umbonatus</i> (Reuss)	-	-	VR	<i>Sigmomorpha</i> spp.	-	VR	-
<i>Oridosalis</i> cf. <i>umbonatus</i> (Reuss)	VR	-	-	<i>Sigmomorpha semitecta terquemiana</i> (Fornasini)	-	-	R
<i>Osangularinella umbonifera</i> (Cushman)	VR	-	-	<i>Sigmomorpha trilocularis</i> (Bagg)	VR	-	-
<i>Paracassidulina sulcata</i> Belford	-	-	R	<i>Sigmovirgulina</i> sp.	-	-	VR
<i>Pararotalia nipponica</i> (Asano)	-	-	R	<i>Sphaeroidina japonica</i> Brady	VR	-	-
<i>Patellina corrugata</i> Williamson	-	VR	R	<i>Sphaeroidina</i> spp.	-	-	-
<i>Patellinella hanzawai</i> Asano	-	-	R	<i>Spirillina limbata</i> Brady	-	-	VR
<i>Pateoris hauerinoides</i> (Rhombler)	-	-	F	<i>Spirillina vivipara</i> Ehrenberg	-	-	R
<i>Planoglabratella australensis</i> (Heron-Allen & Earland)	-	-	R	<i>Spirillina</i> sp.	-	-	VR
<i>Planoglabratella opercularis</i> (d'Orbigny)	-	-	R	<i>Spiroloculina hadai</i> Thalmann	-	-	R
<i>Planoglabratella pateriformis</i> (Brady)	-	-	R	<i>Spiroloculina</i> sp.	-	-	VR
<i>Planoglabratella subopercularis</i> (Asano)	R	R	F	<i>Spirolectammina</i> sp. A	-	-	R
<i>Planoglabratella?</i> sp. C	-	-	VR	<i>Spiropectammina</i> sp.	-	-	VR
<i>Polystomellina discorbinooides</i> Yabe & Hanzawa	-	VR	R	<i>Spirolectinella wrightii</i> (Silvestri)	-	-	R
<i>Poreponides cribrorepondus</i> Asano & Uchio	-	-	R	<i>Textularia conica</i> d'Orbigny	-	-	VR
<i>Pseudodononion japonicum</i> Asano	-	VR	R	<i>Textularia</i> spp.	-	-	R
<i>Pseudodononion</i> spp.	-	VR	VR	<i>Triloculina rotunda</i> d'Orbigny	-	-	VR
<i>Pseudoparrella naraensis</i> Kuwano	VR	-	F	<i>Triloculina suttuensis</i> Asano	-	-	R
<i>Pseudoparrella takayanagi</i> (Iwasa)	VR	R	F	<i>Triloculina tricarinata</i> d'Orbigny	-	-	VR
<i>Pseudoparrella</i> spp.	-	-	R	<i>Triloculina</i> sp. A	-	-	VR
<i>Pseudopolymorpha</i> sp.	-	-	VR	<i>Uvigerina juncea</i> Cushman	-	-	VR
<i>Pullenia apertura</i> Cushman	VR	-	-	<i>Uvigerina</i> spp.	VR	-	-
<i>Pyrgo ezo</i> Asano	VR	-	-	<i>Valvulineria hamanakoensis</i> (Ishiwada)	-	-	VR
<i>Pyrgo fornasinii</i> Chapman & Parr	-	R	-	<i>Valvulineria japonica</i> Asano	-	VR	-
<i>Quinqueloculina agglutinata</i> Cushman	-	-	R	Gen. et sp. indet.	R	F	F
<b>Total counted</b>				<b>869</b>	<b>780</b>	<b>4652</b>	

A: 80.5-13.5%, C: 13.5-4.5%, F: 4.5-1.5%, R: &lt;1.5%, VR: two or less occurrences, -: no occurrence

## 2. Faunal Reference List

Foraminiferal species from the Kuromatsunai and Setana Formations are alphabetically listed below, under the categories of planktons and benthos. The selected species are illustrated with scanning electron micrographs in Plates 1 to 12. The original references are given for species. Additional references are selected from those with helpful remarks and illustrations about the species from the areas studied in this report. All specimens are catalogued and deposited in Department of Science Education, Iwamizawa College, Hokkaido University of Education.

### Planktonic Foraminifera

#### *Globigerina bulloides* d'Orbigny

*Globigerina bulloides* d'Orbigny, 1826, Ann. Sci. Nat., Paris, ser. 1, v. 7, p. 277, Modèle nos. 17, 76; Brady, 1884, Voy. Challenger, Rep., Zool., v. 9, p. 593, pl. 79, figs. 7a-c.

#### *Globigerina quinqueloba* Natland

*Globigerina quinqueloba* Natland, 1938, Bull. Scripps Inst. Oceanogr., Tech. Ser., v. 4, no. 5, p. 149, pl. 6, figs. 7a-c.

#### *Globigerinita glutinata* (Egger)

*Globigerina glutinata* Egger, 1893, Abhandl. K. Bayer. Akad. Wiss. Munchen, CLII, v. 18, p. 371, pl. 13, figs. 19-21.

*Globigerinita glutinata* (Egger), Parker, 1962, Micropal. v. 8, no. 2, p. 246, pl. 9, figs. 1-16.

#### *Globigerinita uvula* (Ehrenberg)

*Pylodexia uvula* Ehrenberg, 1861, K. Preuss., Akad. Wiss. Berlin, Monatsber., p. 276, 277, 308.

*Globigerinita uvula* (Ehrenberg), Parker, 1962, Micropal., v. 8, no. 2, p. 252, pl. 8, figs. 14-26.

#### *Globigerinoides ruber* (d'Orbigny)

*Globigerina rubra* d'Orbigny, 1839, in Sagra, R. de la, Hist. Ohys. Pol. Nat. Cuba, Foraminiferes, p. 82, pl. 4, figs. 12-14.

*Globigerinoides rubra* (d'Orbigny) (sic.), Cushman, 1927, Cushman Lab. Foram. Res., Contr., v. 3, pt. 1, p. 87, pl. 19, figs. 6a-c.

#### *Globorotalia inflata* (d'Orbigny)

*Globigerina inflata* d'Orbigny, 1839, in Barker-Webb and Berthelot, Hist. Nat. Canaries, v. 2, pt. 2, Zool., p. 134, pl. 2, figs. 7-9.

*Globorotalia inflata* (d'Orbigny), sensu strict, Maiya Saito and Sato, 1976, in Takayanagi and Saito eds. Progress in Micropaleontology, Micropal. Press, p. 408, pl. 2, figs. 5-7.

#### *Neogloboquadrina cf. asanoi* (Maiya, Saito, and Sato)

cf. *Globoquadrina asanoi* Maiya, Saito, and Sato, 1976, Progress in Micropaleontology. p.

409, pl. 3, figs. 1a-c, 2a-c, 3.

cf. *Neogloboquadrina asanoi* (Maiya, Saito, and Sato), Thompson, 1980, Init. Rep. DSDP. v. 56-57, pl. 3, figs. 10-12.

*Neogloboquadrina dutertrei* (d'Orbigny)

*Globigerina dutertrei* d'Orbigny, 1839, Foraminiferés, p. 84, pl. 4, figs. 19-21.

*Neogloboquadrina dutertrei* (d'Orbigny), Rögl and Bolli, 1973, Init. Rep. DSDP, v. 27, 743-767, pl. 9, figs. 1-3, 7-10; pl. 17, figs. 1-6.

*Neogloboquadrina incompta* (Cifelli)

*Globigerina incompta* Cifelli, 1961, Cushman Found. Foram. Res., Contr., v. 12, pt. 3, p. 84, pl. 4, figs 1-7.

*Neogloboquadrina incompta* (Cifelli), Rogl and Bolli, 1973, Init. Repts. DSDP, v. 15, p. 571, pl. 10?

*Neogloboquadrina pachyderma* (Ehrenberg)

*Aristerospira pachyderma* Ehrenberg, 1861, K. Preuss. Akad. Wiss, Berlin, Monatsber. p. 276, 277, 303.

*Globigerina pachyderma* (Ehrenberg), Brady, 1884, Voy. Challenger, Rep., Zool., v. 9, p. 600, pl. 114, figs. 19, 20.

*Neogroboquadrina pachyderma* (Ehrenberg), Rogl and Bolli, 1973, Init. Repts. DSDP, v. 15, p. 571, pl. 11, figs. 2-6; pl. 16, fig. 12.

**Benthic Foraminifera**

*Alabamina japonica* (Asano)

*Pseudoparrella japonica* Asano, 1949, Jour. Paleont., 23 (4), p430, fig. 2, Nos. 2-4.

*Alabamina japonica* (Asano), Takayanagi and Hasegawa, 1987, Checklist and bibliography of post-Paleozoic foraminifera established by Japanese workers, 1890-1986, p. 39.

*Ammonia beccarii* (Linné)

*Nautilus beccarii* Linne, 1758, Syst. Nat., se. 10, p. 710.

*Ammonia beccarii* (Linne), Fizzell and Keen, 1949, Jour. Pal., v. 23, no. 1, p. 106.

*Ammonia ketienensis* (Ishizaki)

*Stebius ketienensis* Ishizaki, 1948, Acta Geol. Taiwan, 2 (1), 59, pl. 1, figs. 2a-c.

*Ammonia ketienensis* (Ishizaki), Huang, 1964, Micropaleont. v. 10, no. 1, p. 53, pl. 1, figs. 13a-c.

*Angulogerina hughesi* (Galloway and Wissler)

*Uvigerina hughesi* Galloway and Wissler, 1927, Jour. Pal., v. 1, p.76, pl. 12, figs. 5a, b.

*Angulogerina hughesi* (Galloway and Wissler), Asano, 1950, Pacific Sci. v. 4, no. 2, p. 19, figs. 91-93.

*Angulogerina kokozuraensis* Asano

*Angulogerina kokozuraensis* Asano, 1949, Jour. Pal., v. 23 no. 4, p. 428, text-fig. 1, nos. 50-53; 1950, Illust. Cat.Japan. Tert. Small. Foram., pt. 2, p. 19, figs- 94-96.

*Trifarina kokozuraensis* (Asano), Matoba, 1967, Tohoku Univ., Sci. Rep., 2nd ser. (Geol.) ,

# Foraminifera from the Kuromatsunai and Setana Formations

v.38, no.2, p. 257, pl. 26, fig.3.

*Anomalinella rostrata* (Brady)

*Truncatulina rostrata* Brady, 1881, Quart. Jour. Micr. Sci., n. s., v. 21, p. 65.

*Anomalinella rostrata* (Brady), Cushman, 1927, Cont. Cushman Labo. Foram. Res., v. 3, p. 93.

*Anomalinoides globulosus* (Chapman and Parr)

*Anomalina grosserugosa* Brady, 1884, Challenger Foram., p.675, pl. 94, figs. 4, 5.

*Anomalina globulosa* Chapman and Parr, 1937, Austr. Antarct. Exp. 1911-14, Sci. Rep., ser. C, 1 (2), p.117.

*Astacolus hyalacrulus* Loeblich and Tappan

*Astacolus hyalacrulus* Loeblich and Tappan, 1953, Smithsonian Misc. Coll. v. 121, no. 7, p.52, pl. 9, figs. 1-4.

*Astrononion aomoriense* Asano

*Astrononion aomoriense* Asano, 1950, Illust. Cat. Japan. Tert. Small. Foram., pt.1, p. 5, figs. 25, 26.

*Astrononion hamadaense* Asano

*Astrononion hamadaense* Asano, 1950, Illust. Cat. Japan. Tert. Small. Foram., pt.1, p.6, figs. 29-31.

*Bolivina decussata* Brady

*Bolivina decussata* Brady , 1881, Quart. Jour. Micr. Sci., n.s., v. 21, p. 28; 1884,Voy. Challenger, Rep., Zool., v. 9, p. 423, pl. 53, figs. 12-13.

*Bolivina* sp.A

This form is identified to *Bolivina* sp. A of Hasegawa (1979), and is similar to *Bolivina decussata* Brady in its outline, but differs from it in less developed, and rounded lobes on the wall surface.

*Bolivinita quadrilatera* (Schwager)

*Textularia quadrilatera* Schwager, 1866. Novara Exp. Geol. Theil., 2, p. 253, pl. 7, fig. 10.

*Bolivina quadrilatera* (Schwager), Asano, 1938, Jour. Geol. Soc. Japan, 45, p. 607, pl.16, fig. 10.

*Brizalina alata* (Seguenza)

*Vulvulina alata* Seguenza, 1862, Atti Accad. Gioenia Sci. Nat., 2 (18), p. 115, pl. 2, figs.5, 5a.

*Bolivina alata* (Seguenza), Cushman, 1937, Cushman Lab. Foram., Res. Spac. Publ., 9, p. 106, pl.13, figs. 3-11.

*Brizalina alata* (Seguenza), Belford, 1966, Bureau of Mineral Resources, Geol. and Geophysics, Bull. 79.

*Buccella frigida* (Cushman)

*Pulvinulina frigida* Cushman, 1922, Canada Biol. Contr., no. 9 (1921), p. 12.

*Buccella frigida* (Cushman), Andersen, 1952, Washington Acad. Sci., Jour., v. 42, no. 5, p. 144, figs. 4-6.

*Buccella kuromatsunaiensis* Shirai, 1960, Jour. Fac. Sci. Hokkaido Univ., Ser. IV, vol. X, 537-543, pl. 2, figs. 2a-c.

*Buccella makiyamai* Chiji

*Eponides schreibersii* Morishima and Chiji (not of Reuss), 1952, Mem. Coll. Sci., Univ. Kyoto, ser. B, v. 20, no. 2, pl. 2 (XIII), figs. 6a-c.

*Buccella makiyamae* (sic.) Chiji, 1961, Prof. J. Makiyama, Mem. Vol., Kyoto, p. 234, text-figs. 2a-c, pl. 1, figs. 13-14.

*Buccella nipponica* (Husezima and Maruhasi)

*Discorbis nipponica* Husezima and Maruhasi, 1944, Jour. Shigenkagaku Kenkyusho (Res. Inst. Nat. Resour., Japan), v. 1, no. 3, p. 397, pl. 34. figs. 9a-c.

*Buccella nipponica* (Husezima and Maruhasi), Hasegawa, 1979, Tohoku Univ., Sci. Rep., 2nd ser. (Geol.), v. 49, no. 2, p. 144, pl. 7, figs. 4a-d.

*Buccella tanaii* (Uchio)

*Eponides tanaii* Uchio, 1951, Jour. Geol. Soc. Japan, 57, p. 376, pl. 5, fig. 8.

*Buccella tanaii* (Uchio), Takayanagi and Hasegawa, 1987, Checklist and bibliography of post-Paleozoic foraminifera established by Japanese workers, 1890-1986, p. 19.

*Buccella tenerrima* (Bandy)

*Rotalia tenerrima* Bandy, 1950, Paleont., vol. 24, Tulsa, Okula., p. 278, pl. 42, fig. 3.

*Eponides frigidus* (Cushman), Cushman and Todd, 1947, (not *Pulvinulina frigida* Cushman, 1922), Cushman Lab. Foram. Res. Spec. Publ. v. 23, p. 71, pl. 8, fig. 7.

*Buccella inusitata* Andersen, 1952, Washington Acad. Sci., Jour., v. 42, no. 5, p. 148, figs. 10-11.

*Buccella tenerrima* (Bandy), Feyling-Hanssen, 1976, Maritime Sediments Spec. Publ. 1, p. 353, pl. 1, fig. 8-10; pl. 2, fig. 10-12.

*Bulimina* cf. *imamurae* Tai

cf. *Bulimina imamurae* Tai, 1959, Hiroshima Univ., Jour. Sci. ser. C, v. 2, no. 4, p. 387, pl. 40, fig. 3.

*Buliminella elegantissima* (d'Orbigny)

*Buliminella elegantissima* d'Orbigny, 1839, Voy. Amer. Mérid., Foraminifères, v. 5, pt. 5, p. 51, pl. 7, figs. 13, 14.

*Buliminella elegantissima* (d'Orbigny), Cushman, 1919, U.S. Nat. Mus., Proc., v. 56, p. 606.

*Cancris auriculus* (Fichtel and Moll)

*Nautilus auriculus* Fichtel and Moll, 1789, Testacea microscopica, p. 105, pl. 20, figs. a-f; 1803 var. a, p. 108, pl. 20, figs. a-c; var. b, p. 110, pl. 20, figs. d-f.

*Cancris auricula* (Fichtel and Moll) (sic.), Cushman, 1927, Bull. Scripps Inst. Oceanogr., Tech. Ser., v. 1, no. 10, p. 164, pl. 5. flag. 10.

*Chrysalidinella dimorpha* (Brady)

*Chrysalidina dimorpha* Brady, 1881, Quart. Jour. Micr. Sci., n. s., v. 21, p. 24; 1884, Voy. Challenger, Rep., Zool., v. 9, p. 388, pl. 46, figs. 20-21.

*Chrysalidinella dimorpha* (Brady), Schubert, 1908, Neues Jb. Min. Geol. Pa,1., Beil.-Bd. 25, p. 242.

*Cibicides* cf. *kamadai* Asano

cf. *Cibicides kamadai* Asano, 1951, Illust. Cat. Japan. Tert. Small. Foram., pt. 13, p. 17, figs. 33-35.

*Cibicides lobatulus* (Walker and Jacob)

*Nautilus lobatulus* Walker and Jacob, 1789, Adams Essays, p. 642, pl. 14, fig. 36.

Foraminifera from the Kuromatsunai and Setana Formations

*Cibicides lobatula* (Walker and Jacob) (sic.), Cushman, 1931, U.S. Nat. Mus., Bull. 104, pt. 8, p. 118, pl. 21, figs. 3a-c.

*Cibicides refulgens* de Montfort

*Cibicides refulgens* de Montfort, 1808, Conch. Syst., v. 1, p. 122.

*Truncatulina refulgens* (Montfort), Brady, 1884, Voy. Challenger, Rep., Zool., v. 9, p. 659, pl. 92, figs. 7-9.

*Cibicides subdepressus* (Asano)

*Planulina subdepressa* Asano (sic.), 1951, Illust. Cat. Japan. Tert. Small. Foram., pt. 13, p. 15, figs. 16-18.

*Cibicides subdepressus* (Asano), Takayanagi and Hasegawa, 1987, Checklist and bibliography of post-Paleozoic foraminifera established by Japanese workers, 1890-1986, p. 37.

*Cibicides tani* Iwasa and Kikuchi

*Cibicides tani* Iwasa and Kikuchi, 1954, Paleont. Soc. Japan, Trans. Porc., N. S., no. 16, p. 193, text-figs. 8a-c.

*Cibicidoides pachydermus* (Rzehak)

*Truncatulina pachyderma* Rzehak, 1886, Die Foraminiferenfauna der Neogenformation der Umgebung von Mähr.-Osturf. Naturf. Ver. Brünn, Verh., Bruenn [Brno, Czechoslovakia], vol. 24 (1885), p. 87, pl. 1, fig. 5a-c.

*Truncatulina pseudoungeriana* Cushman, 1922, U.S. Geol. Surv. Prof. Pap., 129 E, p. 97, pl. 20, fig. 9.

*Cibicidoides pachyderma* (sic.), van Morkhoven, 1986, Cenozoic Cosmopolitan Deep-Water Benthic Foraminifera, p. 68-71, pl. 22.

*Cornuspira involvens* (Reuss)

*Operculina involvens* Reuss, 1849, Denkschr. Akad. Wiss., Wien, vol. 1, p. 370, pl. 45, fig. 20.

*Cornuspira involvens* (Reuss), Barker, 1960, SEPM, Spec. Publ., 9, pl. 11, figs. 1-3.

*Cribroelphidium oregonensis* (Cushman and Grant)

*Elphidium oregonense* Cushman and Grant (sic.), 1927, San Diego Soc. Nat. Hist., Trans., V, no. 6, 79, viii, 3.

*Cribroelphidium yabei* (Asano)

*Elphidium yabei* Asano, 1938, Geol. Soc. Japan, Jour., v. 45, no. 538, p. 589, pl. 14, figs. 9-10.

*Cribroelphidium yabei* (Asano), Asano, 1950, Illust. Cat. Japan. Tert. Small. Foram., pt. 1, p. 11, figs. 64-65.

*Cribrolinoides curta* (Cushman)

*Quinqueloculina disparilis* d'Orbigny var. *curta* Cushman, 1917, U.S. Nat. Mus. Bull. v. 71, no. 6, p. 49, pl. 14, fig. 2.

*Cribrolinoides curta* (Cushman), Cusuman and Leroy, 1939, Contr. Cushman Lab. Foram. Res., v. 15, no. 1, p. 17, pl. 3, figs. 1-11; pl. 4, figs. 1-13.

*Cycloforia contrta* d'Orbigny

*Cycloforia contrta* d'Orbigny, 1846, Foram. Foss. Viene, p. 298, pl. 20, figs. 4-6.

*Cystammina pauciloculata* (Brady)

- Trochammina pauciloculata* Brady, 1879, Quarterly Jour. Microscopical Science, new ser. 19, p. 58, pl. 5, figs. 13, 14.
- Anmpchilostoma pauciloculata* (Brady), Eimer and Fickert, 1899, Zeitschrift Wissenschaftliche Zoologie 65, p. 692.
- Cystammina pauciloculata* (Brady), Galloway, 1933, A Manual of Foraminifera. p. 186, pl. 16, fig. 12.
- Discorbinella araucana* (d'Orbigny)
- Rosanila araucana* d'Orbigny, 1839,
- Discorbinella araucana* (d'Orbigny), Jones, 1994, Challenger Foraminifera, p. 93, pl. 86, figs. 10, 11.
- Discorbinella bertheloti* (d'Orbigny)
- Rosalina bertheloti* d'Orbigny, 1839, in Barker-Webb and Berthelot, Hist. nat. Canaries, v. 2, pt. 2, Zool., p. 135, pl. 1, figs. 28-30
- Discorbinella bertheloti* (d'Orbigny), Loeblich and Tappan, 1964, Treatise on Invertebrate Pal., Moore R.C., ed., pt. C, Protista, 2, v. 2, p. C575, figs. 453.
- Dyocibicides perforata* Cushman and Valentine
- Dyocibicides perforata* Cushman and Valentine, 1930, Contr. Stanford Geol. Dept., v. 1, no. 1, p. 31, pl. 10, figs. 3a-c.
- Elphidiella arctica* (Parker and Jones)
- Polystomella arctica* Parker and Jones, 1864, in Brady, Trans. Linn. Soc. London, Zool., v. 24, p. 471, pl. 48, fig. 18.
- Elphidium arcticum* (Parker and Jones), Cushman, 1930, U. S. Nat. Mus. Bull. 104, pt. 7, p. 27, pl. 11, figs. 1-6.
- Elphidiella arctica* (Parker and Jones), Cushman, 1939, U. S. Geol. Surv. Prof. Pap. 191, p. 65, pl. 18, figs. 11-14.
- Elphidium aculeatum* (d'Orbigny)
- Polystomella aculeata* d'Orbigny (sit.), 1846, Foraminifères fossiles du Bassin Tertiare de Vienne (Autriche). Gide et Comp. p. 131, pl. 6, figs. 27-28.
- Elphidium aculeatum* (d'Orbigny), Thalmann, 1932, Eclogae Geologicae Helvetiae, v. 25, 293-312.
- Elphidium advena* (Cushman)
- Polystomella advena* Cushman, 1922, Carnegie Inst. Washington, Publ. 311, p. 56, pl. 9, figs. 11, 12.
- Elphidium advenum* (sit.), Cushman, 1930, U.S. Nat. Mus., Bull. 104, pt. 7, p. 25, pl. 10, figs. 1, 2.
- Elphidium articulatum* (d'Orbigny)
- Polystomella articulata* d'Orbigny, 1839, Amer. Merid., Foraminifères, v. 5, pt. 5, p. 30, pl. 3, figs. 9, 10.
- Elphidium articulatum* (d'Orbigny), Cushman, 1930, U.S. Nat. Mus., Bull. 104, pt. 7, p. 26, pl. 10, figs. 6-8.
- Elphidium bartletti* Cushman
- Elphidium bartletti* Cushman, 1933, Smithsonian Misc. Coll., v. 89, no. 9., p. 4, pl. 1, figs. 9a, b.

Foraminifera from the Kuromatsunai and Setana Formations

*Elphidium bartletti* Cushman, Loeblich and Tappan, 1953, Smithsonian Misc. Coll., v. 121, no. 7, p. 96, pl. 18, figs. 10-14.

*Elphidium crispum* (Linné)

*Nautilus crispum* Linné, 1758, Syst. Nat., ed. 10, p. 709.

*Elphidium crispum* (Linné), Cushman and Grant, 1927, San Diego Soc. Nat. Hist. Trans., v. 5, no. 6, p. 73, pl. 7, figs. 8a, b.

*Elphidium excavatum* (Terquem) forma *clavata* Cushman

*Elphidium incertum* (Williamson) var. *clavatum* Cushman, 1930, U. S. Nat. Mus., Bull. 104, pt. 7, p. 20, pl. 7, figs. 10a, b.

*Elphidium decipiens* (Costa), Hada, 1931, Tohoku Imp. Univ. Sci., Rep., 4th, ser. v. 6, no. 3, p. 126, text-figs. 83a, b.

*Elphidium hughesi foraminosum* Cushman, Asano, 1950, Illust. Cat. Japan. Tert. Small. Foram., pt. 1, p. 8, text-figs. 46, 47.

*Elphidium clavatum* Cushman, Loeblich and Tappan, 1953, Smith. Misc. Coll., v. 121, no. 7, p. 98, pl. 19, figs. 8-10.

*Elphidium excavatum* (Terquem) forma *clavata* Cushman, Takayanagi, 1955, Tohoku Univ., Inst., Geol., Pal., Contr., no. 45, p. 42, pl. 1, fig. 25.

*Elphidium frigidum* Cushman

*Elphidium frigidum* Cushman, 1933, Smithsonian Misc. Coll., vol. 89, no. 9, p. 5, pl. 1, fig. 8.

*Elphidium hanzawai* Asano

*Elphidium hanzawai* Asano, 1939, Jour. Geol., Soc., Japan. v. 46, no 551, 426, figs. 3, 4a, b.

*Elphidium incertum* (Williamson)

*Polystomella umbilicatula* var. *incerta* Williamson, 1858, Recent Foraminifera of Great Britain, p. 44, pl. 3, fig. 82a.

*Elphidium incertum* (Williamson), Macfadyen, 1932, Geol. Mag., vol. 69, no. 821, pl. 35, figs. 6a, b.

*Elphidium jensei* (Cushman)

*Polystomella macella* (Fichtel and Moll) var., Jensen, 1904, Linnean Soc. New South Wales Proc., v. 29, p. 817, pl. 23, fig. 4.

*Polystomella jensei* Cushman, 1924, Carnegie Inst. Washington, Publ. 342, p. 49, pl. 16, figs. 4 (?), 6.

*Elphidium jensei* (Cushman), Cushman, 1933, U.S. Nat. Mus., Bull. 161, pt. 2, p. 48, pl. 11, figs. 6, 7.

*Elphidium macellum* (Fichtel and Moll)

*Elphidium macellus* (Fichtel and Moll) var., 1798, Testacea microscopia, aliaque minuta ex generibus Argonauta et Nautilus, ad naturam picta et descripta. p. 68, pl. 10, figs. h, i, k.

*Elphidium macellum* var. *tumidocamerale* Bogdanowicz, 1932, Oil Geol., Inst., Leningrad, Trans., ser. A, fasc. 22, II, i, 2, text-figs. 9, 10.

*Elphidium macellum* (Fichtel and Moll), Cushman, 1939, Geol. Surv. prof. paper, 191, p. 51, pl. 14, figs. 1-3

*Elphidium subarcticum* Cushman

*Elphidium subarcticum* Cushman, 1944, Cushman Lab. Foram. Res., Spec., Publ., no.

- 12, p. 27. pl. 3, figs. 34, 35.
- Elphidium nakanokawaense* Shirai, 1960, Jour. Fac. Sci. Hokkaido Univ., Ser. IV, vol. X, 537-543, pl. 1, figs. 4a-b, 5a-b.
- Elphidium subgranulosum* Asano
- Elphidium subgranulosum* Asano, 1938, Jour. Geol. Soc. Japan, 45, p. 586, pl. 14, figs. 4a, b.
- Elphidium subincertum* Asano
- Elphidium subincertum* Asano, 1951, Illust. Cat. Japan Tert. Small Foram., pt. 1, p. 10, figs. 56, 57.
- Elphidium translucens* Natland
- Elphidium translucens* Natland, 1938, Scripps Inst. Oceanogr. Bull., Tech. ser., v. 4, p. 144, pl. 5, figs. 3, 4.
- Epistominella pulchella* Husezima and Maruhasi
- Epistominella pulchella* Husezima and Maruhasi, 1944, Jour. Shigenkagaku Kenkyusho (Res. Inst. Nat. Resour., Japan), v. 1, no. 3, p. 398, pl. 34, figs. 10a-c.
- Fissurina annectens* (Burrows and Holland)
- Lagena annectens* Burrows and Holland, 1895, in Jones, Parker and Brady, 1985, Palaeontogr. Soc., London, Part II, p. 203, pt. vii, II.
- Fissurina annectens* (Burrows and Holland), Barker, 1960, Spc. Publ., Soc. Economic Paleontologists and Mineralogists, 9.
- Fissurina baccata* (Hellon-Allen and Earland)
- Lagena orbigniana* var. *baccata* Hellon-Allen and Earland, 1922, Terra Nova Exped., Zool., VI, no. 2, p. 162, pt. vi, figs. 15, 16.
- Fissurina baccata* (Hellon-Allen and Earland), Jones, 1994, Challenger Foram., Pl. 59, Fig. 20.
- Fissurina lacunata* (Burrows and Holland)
- Lagena castrensis* Brady (not of Schwager), 1884, Voy. Challenger, Rep. Zool., v. 9, p. 485, pl. 60, figs. 1, 2.
- Lagena lacunata* Burrows and Holland, 1895, in Jones, T.R., Palaeontogr. Soc. London, p. 205, pl. 7, fig. 12.
- Fissuriua orbigniana lacunata* (Burrows and Holland), Asano, 1938, Tohoku Imp. Univ., Sci. Rep., 2nd ser. (Geol.), v. 19, no. 2, p. 219, pl. 27, figs. 27, 28.
- Fissurina lucida* (Williamson)
- Entosolenia marginata* (Montagu) var. *lucida* Williamson, 1848, Ann. Mag. Nat. Hist., ser. 2, v. 1, p. 17, pl. 2, fig. 17.
- Fissurina lucida* (Williamson), Loeblich and Tappan, 1953, Smithsonian Misc. Coll., v. 121, no. 7, p. 76, pl. 14, fig. 4.
- Fissurina marginata* (Montagu)
- Vermiculum marginatum* Montagu, 1803, Testacea Britannica, p. 524.
- Fissurina marginata* (Montagu), Loeblich and Tappan, 1953, Smithsonian Misc. Coll., v. 121, no. 7, p. 77, pl. 14, figs. 6-9.
- Fissurina obscurocostata* Galloway and Wissler
- Fissurina obscurocostata* Galloway and Wissler, 1927, Jour. Pal., v. 1, p. 52, pl. 9, fig. 1.

Foraminifera from the Kuromatsunai and Setana Formations

*Fissurina orbignyana* Seguenza

*Fissurina orbignyana* Seguenza , 1862, Foram. monotal. Mioc. Messina, p. 66, pl. 2, figs. 24, 26.

*Fissurina cf. rizzae* Seguenza

cf. *Fissurina rizzae* Seguenza, 1862, Foram. monotal., Mioc. Messina, p. 72, pl. 2, fig. 50.

*Fissurina semimarginata* (Reuss)

*Lagena* sp. (Nos. 64-65) von Schlicht, 1870. Die Foraminiferen Septarienthones Pielzpuhl, p. 11, pl. 4, figs. 4-6, 10-12.

*Lagena marginata* Williamson var. *semimarginata* Reuss, 1870, Sitzb. Akad. Wiss. Wien, v. 62, pt. 1, p. 468.

*Fissurina semimarginata* (Reuss), Barker, 1960, Spc.Publ., Soc. Economic Paleontologists and Mineralogists, 9.

*Fissurina cf. subquadrata* Parr

cf. *Fissurina subquadrata* Parr , 1954, Roy. Soc. Victoria, Proc., n. s., LVI, pt. 2, 203, ix, 5.

*Gavelinopsis praegeri* (Heron-Allen and Earland)

*Discorbina praegeri* Heron-Allen and Earland, 1913, Royal Irish Acad., Proc., v. 31, pt. 64, p. 122, pl. 10, figs. 8-10.

*Gavelleopsis praegeri* (Heron-Ailen and Earland), Hofker, 1951, Siboga Exped., Mon. IV, pt. 3, p. 485. figs. 332-334.

*Glabratella cf. aurantista* Seiglie and Bermúdez

cf.*Glabratella aurantista* Seiglie and Bermúdez, 1965, Monographia de la familia de foraminíferos Glabratellidae, Geos, 1965, no. 12,

*Glabratella mirabiris panamensis* Seiglie and Bermúdez

*Glabratella mirabiris panamensis* Seiglie and Bermúdez , 1965, Geos, 12, p. 33, pl. 6, a-b, 7, 8.

*Glabratella pulvinata* (Brady)

*Discorbina pulvinata* Brady, 1884, Rept. on the Sci. Reslt. of the Voy. of the H. M. S. Challenger during the years 1873-1876, Zoology, v. 9,

*Glabratella pulvinata*, (Brady), Barker, 1960, SEPM Spec. Publ., 9, pl. 88, figs. 10a, b.

*Glabratella* sp.A

This species is somewhat resembles *G. crassa* Hofker but differs from it in having more rounded and inflated chambers, and five to six chambers in final whorl.

*Glabratella* sp.B

This species is characterized in having planoconvex test with low trochospiral coiling and with long spines in its periphery.

*Glandulina ovula* d'Orbigny

*Glandulina ovula* d'Orbigny, 1846, Foram. Foss. Bass. Tert. Vienne, p. 29, pl. 1, figs. 6, 7.

*Globocassidulina bisecta* Nomura

*Globocassidulina bisecta* Nomura, 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 53, no. 1, p. 73-77, pl. 2,figs. 2,3; pl. 14, figs. 8-12; pl. 15, figs. 1-5.

*Globocassidulina jamesoni* (McCulloch)

*Cassidulina* (?) *jamesoni* McCulloch, 1977, Quantitative obserbations on Recent foram. tests, with emphasis on the eastern Pacific, p. 390, pl. 164, figs. 15a-c.

*Globocassidulina jamesoni* (McCulloch), Nomura, 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 54, no. 1, p. 33, 34, pl. 3, figs. 7a-c, 8a, b; pl. 19, figs. 1, 2.

*Globocassidulina canalisuturata* Eade

*Globocassidulina canalisuturata* Eade, 1967, N.Z. Jour. Mar. Freshwater Res., 1. no. 4, p. 440, fig. 3 (5-7), fig. 5 (7, 8).

*Globocassidulina neobrecha* Nomura

*Globocassidulina neobrecha* Nomura, 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 53, no. 1, p. 67-68, pl. 1, figs. 16a-c; pl. 19, figs. 3-6.

*Guttulina yabei* Cushman and Ozawa

*Guttulina yabei* Cushman and Ozawa, 1929, Japan Jour. Geol. Geogr., v. 6, nos. 3-4, p. 68, pl. 13, fig. 2, pl. 14, fig. 6.

*Hanzawaia nipponica* Asano

*Hanzawaia nipponica* Asano, 1943, Jour. Geol. Soc. Japan, v. 51, no. 606, p. 98, pl. 4, figs. 1-2.

*Heterolepa praecincta* (Karrer)

*Rotalia praecinctus* Karrer, 1868, Sitz. Akad. Wiss. Wien, v. 58, p. 189, pl. 5, fig. 7.

*Truncatulina praecincta* Cusuman, 1915, U.S. Nat. Mus. Bull., v. 71, no. 5, p. 39, pl. 26, fig. 2. *Eponides praecinctus* (Karrer), Asano, 1951, Illust. Cat. Japan Tert. Small Foram., pt. 14, p. 11, figs. 30-32.

*Heterolepa subhaidingerii* (Parr)

*Cibicides subhaidingerii* Parr, 1950, Foraminifera, Rep. B.A.N.Z. Antarctic Rese. Exp. 1929-1931, Ser. B. (Zoology and Bontany), 5, p. 364, pl. 15, fig. 7.

*Heterolepa subhaidingerii* (Parr), Tappan and Loeblich, 1982, Synopsis and Classification Living Organisms, pl. 53, fig. 10.

*Islandiella helenae* Feyling-Hanssen and Buzas

*Cassidulina teretis* Tappan, Loeblich and Tappan (not of Tappan, 1951), 1953, Smithsonian Misc. Coll., v. 121, no. 7, p. 121, pl. 24, figs. 3, 4.

*Islandiella helenae* Feyling-Hanssen and Buzas, 1976, Jour. Foram. Res., v. 6, no. 2, p. 155, figs. 1-4.

*Islandiella islandica* (Nørbang)

*Cassidulina islandica* Nørbang, 1945, Foraminifera. in the Zoology of Iceland. v.2, p. 42, text-figs. 7, 8d-f.

*Islandiella islandica* (Nørbang), Nørbang, 1958, Vidensk. Medd. Dansk Natur-hist. Foren., v. 120, p. 27, 28, pl. 6, figs. 1a, b, 2-5, pl. 7, figs. 6a,b, 7a, b,

*Islandiella japonica* (Asano and Nakamura)

*Cassidulina japonica* Asano and Nakamura, 1937, Japan Jour. Geol. Geogr., v. 14, nos. 3-4, p. 144, pl. 13, figs. 1, 2.

*Islandiella japonica* (Asano and Nakamura), Troitskaja, 1970, Usloviya obitaniya i raspredelenie foraminifer v Yaponskom More (semeistva Elphidiidae, Cassidulinidae i Islandiellidae). In Fursenko A.V. ed., Obstchie voprosy izucheniya mikrofauny Sibiri, dalinego vostoka i drugikh raionov. Instituta Geologii i Geofiziki, Akademiya Nauk SSSR, Sibirskoe Otdelenie, Trudy, vol. 71, p.150, pl. 6, figs. 3,4.

*Islandidlla norcrossi* (Cushman)

Foraminifera from the Kuromatsunai and Setana Formations

*Cassidulina norcrossi* Cnshman, 1933, Smithsonian Misc. Coll., v. 89, no. 9, p. 7, pl. 2, figs. 7a-c.

*Islandiella norcrossi* (Cushman), Nørbang, 1958, Dansk Naturh. Foren. Kobenhavn Vidensk. Meddel., v. 120, p.26.

*Islandiella setanaensis* (Asano and Nakamura)

*Cassidulina setanaensis* Asano and Nakamura, 1937, Japan Jour. Geol. Geogr., v. 14, nos. 3-4, pl. 13, figs. 7a,b

*Islandiella setanaensis* (Asano and Nakamura) , Nomura, 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 53, no. 1, p. 1-101, pl. 3, fig. 6, pl.4, figs. 3,4, pl. 10, fig. 11, pl.11, fig. 1-3; 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 54, no. 1, p. 5, pl. 1, figs. 3a-c.

*Islandiella sublimbata* (Asano and Nakamura)

*Cassidulina sublimbata* Asano and Nakamura, 1937, Japan Jour. Geol Geogr., v. 14, nos. 3-4, p. 146, pl. 14, figs. 3, 4a, b.

*Islandiella sublimbata* (Asano and Nakamura), Nomura, 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 53, no. 1, pl. 3, fig. 7, pl. 4, fig. 2, pl. 5, fig. 7, pl. 9, figs. 1-8; 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 54, no. 1, p. 9, pl. 1, figs. 6-8.

*Islandiella yabei* (Asano and Nakamura)

*Cassidulina yabei* Asano and Nakamura. 1937, Japan. Jour. Geol. Geogr., v. 14, nos. 3-4, p. 145, pl. 14, figs. 1a, b.

*Islandiella yabei* (Asano and Nakamura), Nomura, 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 53, no. 1, pl. 3, fig. 1; pl. 4, fig. 7; pl. 9, figs. 9-12, pl. 10, figs. 1-3.

*Karreriella baccata* (Schwager) *japonica* Asano

*Karreriella baccata* (Schwager) *japonica* Asano, 1938, Japanese Jour. Geol. Geogr., v.15, nos. 1-2, p. 90, pl. 10, figs. 1a-c.

*Lagena acuticosta* Reuss

*Lagena acuticosta* Reuss, 1861, Sitz. Akad. Wiss. Wier., v. 44, no. 1, p. 305, pl. 1, fig. 4.

*Lagena apiopleura* Loeblich and Tappan

*Lagena apiopleura* Loeblich and Tappan, 1953, Smithsonian Misc. Coll., v. 121, no. 7, p. 59, pl. 10, figs. 14, 15.

*Lagena striata* (d'Orbigny)

*Oolina striata* d'Orbigny, 1839, Voy. Amer. Merid., Foraminifères, v. 5, pt. 5, p. 21, pl. 5, fig. 12.

*Lagena striata* (d'Orbigny), Asano, 1938, Tohoku Imp. Univ., Sci. Rep., 2nd ser. (Geol.), v. 19, p. 217, pl. 27, fig. 26; pl. 28, fig. 28.

*Lagena sulcata spicata* Cushman and McCulloch

*Lagena sulcata* var. *spicata* Cushman and McCulloch, 1950, Allan Hancock Pacific Exped., v. 6, no. 6, p. 360, pl. 48, figs. 3-7.

*Lenticulina nikobarensis* (Schwager)

*Cristellaria nikobarensis* Schwager, 1886, Novara Exp. Geol. Thell., 2, p. 243, pl. 6, fig. 87.

*Robulus nikobarensis* (Schwager), Asano, 1938, Sci. Rep. Tohoku Univ. ser. 2, 19 (2), p. 204, pl. 28, figs. 5, 6; pl. 29, fig. 8.

*Lenticulina nikobarensis* (Schwager), Loeblich and Tappan, 1988, Foraminiferal genera and their classification. p. 405. Pl. 446, figs. 9, 10.

*Melonis pompilioides* (Fichtel and Moll)

*Nautilus pompilioides* Fichtel and Moll, 1798, Testacea microscopica, p. 31, pl. 2, figs. a-c.

*Melonis etruscus* Montfort, 1808, Conch. Syst., p. 67.

*Melonis pompilioides* (Fichtel and Moll), Voloshinova, 1958, Mikrofauna SSSR, Sbornik 9,

VNIGRI, Trudy, p. 149, pl. 3, fig. 1

*Melonis uchiori* Hasegawa

*Melonis uchiori* Hasegawa, 1991, Trans. Proc. Palaeont. Soc. Japan, N.S., no. 164, p. 1003.

*Melonis pacificus* (Cushman)

*Nonion umbilicatula* var. *pacifica* Cushman, 1924, Carnegie Inst. Wash., v. 342, p. 48, pl. 16, fig. 3.

*Nonion pacificum* Cushman, 1939, U.S. Geol. Surv., Prof. paper v. 191, p. 25, pl. 6, fig. 25.

*Melonis pacificus* (Cushman), Hasegawa, 1979, Tohoku Univ., Sci. Rep., 2nd ser. (Geol.), v. 49, no. 2, p. 151.

*Miliolinella circularis* (Bornemann)

*Triloculina circularis* Bornemann, 1855, Zeit. deutsch. Geol. Ces., v. 7, p. 349, pl. 19, fig. 4

*Miliolinella circularis* (Bornemann), Asano, 1951, Illust. Cat. Japan. Tert. Small. Foram., pt. 6, p. 9, figs. 65-67.

*Neoconorbina stachi* (Asano)

*Discopulvinulina stachi* Asano, 1951, Illust. Cat. Japan Tert. Small. Foram., pt. 14, p. 7, figs. 46-48.

*Neoconorbina stachi* (Asano), Matoba, 1970, Tohoku Univ., Sci. Rep., 2nd ser. (Geol.), v. 42, no. 2, p. 57, pl. 4, figs. 6a-c.

*Neoeponides procerus* (Brady)

*Pulvinulina procura* Brady, 1884, Voy. Challenger, Rep., Zool., v. 9, p. 698, pl. 105, figs. 7a-c. *Neoeponides procerus* (Brady), Reiss, 1960, Israel Geol. Surv., Bull. no. 29, p. 17.

*Nonion japonicus* Asano

*Nonion japonicum* Asano (sic.), 1938, Jour. Geol. Soc. Japan, v. 45, no. 593, pl. 15(4), figs. 1, 2.

*Nonionella stella* Cushman and Moyer

*Nonionella miocenica* Cushman var. *stella* Cushman and Moyer, 1930, Cushman Lab. Foram. Res., Contr., v. 6, pt. 1, p. 56, pl. 7, figs. 17a-c.

*Nonionellina labradorica* (Dawson)

*Nonionina labradorica* Dawson, 1860, Canad. Nat., v. 5, p. 191, text-fig. 4.

*Nonionellina labradorica* (Dawson), Voloshinova, 1958, Mikrofauna SSSR, Sbornik 9, VNIGRI, Trudy, no. 115, p. 142.

*Nonionoides grateloupi* (d'Orbigny)

*Nonionina grateloupi* d'Orbigny, 1826, Annales des Sciences Naturelles v. 7, p. 294,

*Nonionoides grateloupi* (d'Orbigny), Saidova, 1975, Bentosnye Foraminifery Tikhogo Okeana, 3 vol., p. 248.

*Oolina carteri* Albani and Yassini

*Oolina carteri* Albani and Yassini, 1989, Aust. J. Mar. Freshw. Res., v. 70, p. 385, fig. 3r-s.

*Oolina costata* (Williamson)

*Entosolenia costata* Williamson, 1858, Recent Foraminifera Great Britain, p. 9, pl. 1, fig.

18.

*Lagena costata* (Williamson), Cushman, 1923, U.S. Nat. Mus. Bull. v. 104, pt. 4, p. 12, pl. 1, fig. 16, pl. 2, figs. 1, 2 (not pl. 3, fig. 8)

*Oolina costata* (Williamson), Parker, 1952, Bull. Mus. Comp. Zool., v. 106, no. 9, p. 409, pl. 4, figs. 20, 21.

*Oolina melo* d'Orbigny

*Oolina melo* d'Orbigny, 1839, Voy. Amer. Merid., Foraminifères, v. 5, pt. 5, p. 20, pl. 5, fig. 9.

*Oolina striatopunctata* (Parker and Jones)

*Lagena sulcata* (Walker and Jacob) var. *striatopunctata* Parker and Jones, 1865, Philos. Trans. Roy. Soc. London, v. 155, p. 350, pl. 13, figs. 25-27.

*Entosolenia striatopunctata* (Parker and Jones) Dawson, 1870, Can. Nat., n.s., v. 5, p. 178, fig. 11. *Lagena striatopunctata* Parker and Jones, Brady, 1878, Ann. Mag. Nat. Hist., ser. 5, v. 1, p. 434, pl. 20, fig. 3.

*Oolina striatopunctata* (Parker and Jones) Loeblich and Tappan, 1953, Smithsonian Misc. Coll. v. 121, no. 7, p. 74, pl. 12, figs. 2-5.

*Oridorsalis umbonatus* (Reuss)

*Rotalia umbonata* Reuss, 1851, Deutsch. geol. Gesell., Zeischr., v. 3, p. 75, pl. 5, figs. 35a-c.

*Oridorsalis umbonatus* (Reuss), Parker, 1964, Jour. Pal., v. 38, no. 4, p. 626, pl. 99, figs. 4-6.

*Osangulariella umbonifera* (Cushman)

*Pulvinulinella umbonifera* Cushman, 1933, Cushman Lab. Foram. Res., Contr., 9, 90.

*Osangulariella umbonifera* (Cushman), Jones, 1994, Challenger Foraminifera, p. 99, pl. 95, figs. 9, 10.

*Paracassidulina sulcata* Belford

*Paracassidulina sulcata* Belford, 1966, Bur. Min. Resour. Aust. Rep., no. 79, p. 142, 144, pl. 24, figs. 11-14, text-fig. 16, nos. 7, 8.

*Pararotalia nipponica* (Asano)

*Rotalia nipponica* Asano, 1936, Jour. Soc. Japan, v. 43, no. 515, p. 614, pl. 30, figs. 2a-c.

*Pararotalia nipponica* (Asano), Ujiie, 1966, Palaeont. Soc. Japan, Trans. Proc., N.S.m no. 61, p. 192, text-figs. 1-3; pl. 24, figs. 1-7; pl. 25, figs. 1-5.

*Patellina corrugata* Williamson

*Patellina corrugata* Williamson, 1858, Rec. Foram. Gt. Britain, p. 46, pl. 3, figs. 86-89.

*Patellinella hanzawai* Asano

*Patellinella hanzawai* Asano, 1936, Jour. Geol. Soc. Japan, v. 43, no. 515, p. 613, pl. 31, figs. 3a-c.

*Pateoris hauerinoides* (Rhumbler)

*Quinqueloculina subrotunda* (Montagu) forma *hauerinoides* Rhumbler, 1936, Foram. der Kieler Bucht, Teil II-Ammodisculinidae bis Textulinidae, v. 1, no. 1, p. 206, 217, 226, text-figs. 167, 208-212.

*Pateoris hauerinoides* (Rhumbler) Loeblich and Tappan, 1953, Smithsonian, Misc. Coll. v. 121, no. 7, p. 42, pl. 6, figs. 8-12, text-figs. 1A, B.

*Planoglabratella australensis* (Heron-Allen and Earland)

*Discorbis australensis* Hellon-Allen and Earland, 1932, Discovery Repts., IV, 416.

*Glabratella australensis* (Heron-Allen and Earland), Loeblich and Tappan, 1964, Treatise on Invertebrate. p. 588.

*Planoglabratella opercularis* (d'Orbigny)

*Rosalina opercularis* d'Orbigny, 1839, Foraminifères, p. 93, pl. 3, figs. 24, 25.

*Discorbis opercularis* Cushman, 1915, U.S. Nat. Mus. Bull., vol. 71, no. 5, p. 18, pl. 11, fig. 3.

*Planolabratella opercularis* (d'Orbigny), Seiglie and Bermúdez, 1965, Geos, 1965, no. 12, 15-65.

*Planoglabratella patelliformis* (Brady)

*Discorbina patelliformis* Brady, 1884, Rept. on the Sci. Reslt. of the Voy. of the H. M. S. Challenger during the years 1873-1876, Zoology, v. 9,

*Glabratella patelliformis* (Brady), Seiglie and Bermúdez, 1965, Monographia de la familia de foraminíferos Glabratellidae, Geos, 1965, no. 12,

*Planoglabratella subopercularis* (Asano)

*Discorbis subopercularis* Asano, 1951, Illust. Cat. Japan Tert. Small Foram., pt. 14, p. 3, figs. 17-19.

*Glabratella subopercularis* (Asano), Matoba, 1970, Tohoku Univ., Sci. Rep., 2nd ser. (Geol.), v. 42, no. 1, p. 54, pl. 5, figs. Sa-c.

*Planolabratella subopercularis* (d'Orbigny), Seiglie and Bermúdez, 1965, Geos, 1965, no. 12, 15-65.

*Planoglabratella* ? sp. C

This species resembles *Planoglabratella subopercularis* (Asano). but differs from it in havng less number of chambers in final whorl.

*Polystomelina discorbinoides* Yabe and Hanzawa

*Polystomelina discorbinoides* Yabe and Hanzawa , 1923, Japan Jour. Geol. Geogr., v. 2, p. 99, text-flgs. a-c.

*Poroeponides cribrorepandus* Asano and Uchio

*Poroeponides cribrorepandus* Asano and Uchio, 1951, Illust. Cat. Japan. Tert. Small. Foram., pt. 14, p.18, figs. 134, 135.

*Pseudononion japonicum* Asano

*Pseudononion japonicum* Asano, 1936, Jour. Geol. Soc. Japan, v. 43, no. 512, p. 347, text-flgs. a-c.

*Pseudoparella naraensis* Kuwano

*Pseudoparella naraensis* Kuwano, 1950, Jour. Geol. Soc. Japan, v. 56, no. 657, p. 317, text-flgs. 6a-c.

*Pseudoparella takayanagii* (Iwasa)

*Epistominella takayanagii* Iwasa, 1955, Jour. Geol. Soc. Japan, v. 61, no. 712, p. 16, text-flgs. 4a-c.

*Epistominella suttsuensis* Shirai, 1960, Jour. Fac. Sci. Hokkaido Univ., Ser. IV, vol X, 537-543, pl. 2,figs. 3a-c.

*Pseudoparella takayanagii* (Iwasa), Hasegawa, 1979, Tohoku Univ., Sci. Rep., 2nd ser.

Foraminifera from the Kuromatsunai and Setana Formations

(Geol.), v. 49, no. 2, p. 153, pl. 5, figs. 7a-c.

*Pseudopolymorphina suboblonga* Cushman and Ozawa

*Pseudopolymorphina suboblonga* Cushman and Ozawa, 1930, U.S. Nat. Mus., Proc., v. 77, p. 91, pl. 23, figs. 3a-c.

*Pullenia apertura* Cushman

*Pullenia apertura* Asano (not of Cushman, 1927), 1951, Illust. Cat. Japan. Tert. Small. Foram., pt. 12, p. 10, figs. 3, 4.

*Pyrgo ezo* Asano

*Pyrgo ezo* Asano, 1938, Japanese Jour. Geol. Geogr., v. 15, nos. 1-2, p. 93, pl. 9, figs. 1-4, 6.

*Pyrgo fomasinii* Chapman and Parr

*Pyrgo fomasinii* Chapman and Parr, 1935, Roy. Soc. W. Australia, Jour., XXI, p. 5.

*Quinqueloculina agglutinata* Cushman

*Quinqueloculina agglutinata* Cushman, 1917, U.S. Nat. Mus., Bull. 71, pt. 6, p. 43, pl. 9, figs. 2a-c.

*Quinqueloculina akneriana* d'Orbigny

*Quinqueloculina akneriana* d'Orbigny, 1846, Foram. Foss. Bas. Vienne, p. 290, pl. 18, figs. 16-21.

*Quinqueloculina costata* d'Orbigny

*Quinqueloculina costata* d'Orbigny, 1826, Ann. Sci. Nat., v. 7, p. 301. no. 3.

*Quinqueloculina elongata* Natland

*Quinqueloculina elongata* Natland, 1938, Bull. Scripps Inst. Oceanogr. Tech. Ser., v. 4, no. 5, p. 141, pl. 4, fig. 5.

*Quinqueloculina kuromatunaiensis* Asano

*Quinqueloculina kuromatunaiensis* Asano, 1936, Jour. Geol. Soc. Japan, v. 43, no. 515, p. 621, pl. 32, figs. 4a-d.

*Quinqueloculina sawanensis* Asano

*Quinqueloculina sawanensis* Asano, 1951, Illust. Cat. Japan. Tert. Small. Foram., pt. 6, p. 6, figs. 40-42.

*Quinqueloculina seminulum* (Linné)

*Serpula seminulum* Linné, 1758, Syst. Nat., ed. 10, p. 786.

*Quinqueloculina seminulum* (Linné), d'Orbigny, 1826, Ann. Sci. Nat., Paris, ser. I, v. 7, . 303, no. 4.

*Quinqueloculina vulgaris* d'Orbigny

*Quinqueloculina vulgaris* d'Orbigny, 1826, Ann. Sci. Nat., v. 7, p. 302, no. 33.

*Quinqueloculina cf. vulgaris* d'Orbigny

cf. *Quinqueloculina vulgaris* d'Orbigny , 1826, Ann. Sci. Nat., v. 7, p. 302, no. 33.

*Quinqueloculina yessoensis* Asano

*Quinqueloculina yessoensis* Asano, 1936, Jour. Geol. Soc. Japan. 43, p. 620, pl. 32, figs. 3a-c.

*Quinqueloculina* sp.B

This species is somewhat similar to *Quinqueloculina elongata* Natland of Asano (1950) in its outline, but is distinguished from it by more inflated chambers and coarsely perforated wall.

*Quinqueloculina* sp.C

This species is somewhat similar to *Quinqueloculina contrita* d'Orbigny, but is distinguished by more rounded outline, short size, and apertural view.

*Quinqueloculina* sp.F

This form is similar to *Quinqueloculina yessoensis* Asano, but is differs from it in having a broadly oval test with more sharply angled periphery in apertural view. This is also similar to *Quinqueloculina kuromatunaiensis* Asano, but is distinguished in having angled periphery of chambers.

*Quinqueloculina* sp.G

This form is somewhat similar to *Quinqueloculina elongata* Natland, but is distinguished from it by its triangular appearance in apertural view.

*Quinqueloculina* sp.H

This form is similar to *Quinqueloculina hasimotoi* Asano, but differs from it in having a test without apertural protrusion.

*Rectobolivina raphanuss* (Parker and Jones)

*Uvigerina (Sagrina) raphana* Parker and Jones, 1865, Roy. Soc. London Philos. Trans., v. 155, p. 364, pl. 18, figs. 16, 17.

*Rectobolivina raphana* (sic.) (Parker and Jones), Loeblich and Tappan, 1964, Treatise on Invertebrate Pal., Moore, R.C., ed., pt. C, Protista 2, v. 2, p. C533, figs. 438 (9-11).

*Rosalina australis* (Parr)

*Discorbis australis* Parr, 1932, Roy. Soc. Victoria, Proc., XLIV (n.s.) p. 227, pl. xxii, 31.

*Rosalina australis* (Parr), Jones, 1994, Challenger Foram., pl. 87, figs. 5-7.

*Rosalina bradyi* (Cushman)

*Discorbina globularis* Brady (not of d'Orbigny, 1826), 1884, Voy. Challenger, Rep., Zool., v. 9, p. 86, figs. 8a-c.

*Discorbina globularis* var.*bradyi* Cushman, 1915, U.S. Nat. Mus., Bull. 71, pt. 5, p. 12, pl. 8, figs. 1a-c.

*Rosalina bradyi* (Cushman), Hornbrook and Vella, 1954, Micropal., v. 8, no. 1, p. 26.

*Rosalina globularis* d'Orbigny

*Rosalina globularis* d'Orbigny, 1926, Ann. Sci. Nat., Paris, ser. 1, v. 7, p. 271, no. 1, pl. 13, figs. 1-14; Modele no. 69.

*Rosalina isabelleana* d'Orbigny

*Rosalina isabelleana* d'Orbigny, 1839, Foraminifères. p. 43, pl. 6, figs. 10-12.

*Rosalina vilardeboana* d'Orbigny

*Rosalina vilardeboana* d'Orbigny, 1839, Voy. Amer. Merid., Foraminifères, v. 5, pt. 5, p. 44, pl. 6, figs. 13-15.

*Rosalina vilardeboana* d'Orbigny

*Rosalina vilardeboana* d'Orbigny, 1839, Voy. Amér. Mérid., Foraminifères, v. 5, pt. 5, p. 44, pl. 6, figs. 13-15.

*Sigmoidella pacifica* Cushman and Ozawa

*Sigmoidella (Sigmoidina) pacifica* Cushman and Ozawa, 1928, Cushman Lab. Foram. Res., Contr., v. 4, p. 19, pl. 2, fig. 13.

*Sigmoilina sigmoidea* (Brady) *compressa* Cushman

Foraminifera from the Kuromatsunai and Setana Formations

- Sigmoilina sigmoidea* (Brady) *compressa* Cushman, 1946, Contr. Cushman Lab. Foram. Res., 22, no. 2, p. 32, pl. 5, figs. 10-12.
- Sigmomorphina semitecta terquemiana* (Fornasini)  
*Polymorphina amygdaloides* Reuss var. *terquemiana* Fornasini, 1902, Mem. Accad. Inst. Sci. Bologna, ser. 5, no. 9, p. 72, fig. 25.  
*Sigmomorphina semitecta* (Reuss) var. *terquemiana* Cushman and Ozawa, 1930, U.S. Nat. Mus. Proc., v. 77, no. 6, p. 129, pl. 33, figs. 4, 5; pl. 34, figs. 2, 3; pl. 35, fig. 1.
- Sigmomorphina trilocularis* (Bagg)  
*Polymorphina trilocularis* Bagg, 1912, U. S. Geol. Surv. Bull., 513, p. 75, pl. 20, figs. 15-18.  
*Sigmomorphina trilocularis* (Bagg), Cushman and Ozawa, 1930, U. S. Nat. Mus. Proc., v. 77, no. 6, p. 136, pl. 36, fig. 5.
- Sphaeroidina japonica* Asano  
*Sphaeroidina japonica* Asano, 1953, Tohoku Univ., Inst. Geol. Paleont., Short Papers, no. 5, p. 17, pl. 2, figs. 43, 44.
- Spirillina limbata* Brady  
*Spirillina limbata* Brady, 1879, Quart. Jour. Microscopic Sci., n.s., vol. 19, p. 64, pl. 8, figs. 26a, b.
- Spirillina vivipara* Ehrenberg  
*Spirillina vivipara* Ehrenberg, 1843, Phys. Ab. K. Akad. Wiss. Berlin, 1841 (1), p. 323, 422, pl. 3, VII, fig. 41.
- Spiroloculina hadai* Thalmann  
*Spiroloculina costata* Hada, 1931, Tohoku Univ. Sci. Rep., 4th ser., Biol., v. 6, no. 1, p. 84, text-figs. 37a,b.  
*Spiroloculina hadai* Thalmann, 1933, Jour. Paleont., v. 7, no. 30, p. 354.
- Spiroplectammina* sp.A  
This form somewhat similar to *Textularia candéiana* d'Orbigny, but has a test with planispirally arranged chambers in the early portion.
- Spiroplectinella wrightii* (Silvestri)  
*Spiroplecta wrightii* Silvestri, 1903, Accad. Pont. Nuovi Lincei, Atti, LVI, p. 59, pl. 1x, figs. 1-6.
- Textularia conica* d'Orbigny  
*Textularia conica* d'Orbigny, 1839, Foraminifères, p. 143, pl. 1. figs. 19, 18.
- Triloculina rotunda* d'Orbigny  
*Triloculina rotunda* d'Orbigny, 1826, Ann. Sci. Nat. v. 7, p. 299, no. 4.
- Triloculina suttensis* Asano  
*Triloculina suttensis* Asano, 1936, Jour. Geol. Soc. Japan, v. 43, no. 515, p. 621, pl. 33, figs. 2a-c.
- Triloculina tricarinata* d'Orbigny  
*Triloculina tricarinata* d'Orbigny, 1826, Ann. Sci. Nat., 7, p. 299, no. 7.
- Uvigerina akitaensis* Asano  
*Uvigerina akitaensis* Asano, 1950, Illust. Cat. Japan. Tert. Small. Foram. pt. 2, p. 14, figs 60-62.

*Uvigerina juncea* Cushman and Todd

*Uvigerina juncea* Cushman and Todd, 1941, Cushman Lab. Foram. Res., Contr., XVII, p. 78, pl. xx, 4-11.

*Uvigerina schencki* Asano

*Uvigerina schencki* Asano, 1950, Illust. Cat. Japan. Tert. Small. Foram. pt. 2, p. 17, figs. 74, 75.

*Valvularia hamanakoensis* (Ishiwada)

*Anomalina hamanakoensis* Ishiwada, 1958, Geol. Surv. Japan Rep., no. 180, p. 18, text-figs. 3a-c; pl. 1, 24-27.

*Valvularia hamanakoensis* (Ishiwada), Matoba, 1970, Tohoku Univ., Sci. Rep., 2nd ser. (Geol.), v. 42, no. 1, p. 63, pl. 4, figs. 12, 13.

*Valvularia japonica* Asano

*Valvularia japonica* Asano, 1951, Illust. Cat. Japan. Tert. Small. Foram. pt. 14, p. 7, figs. 49-51.

## Acknowledgments

The author would like to express his deep gratitude to Associate Professor Shiro Hasegawa of the Laboratory of Geosphere Science, Graduate School of Environmental Earth Science, Hokkaido University, for his encouragement and advice during the course of this study. The author also wishes to thank Professor Yoshihiro Togo and Associate Professor Akihiko Suzuki of the Department of Earth Science, Iwamizawa College, Hokkaido University of Education, for their suggestions concerning various aspects of the present study. Sincere thanks are due to Dr. Yoshinori Hikida of Nakagawa Museum, Mr. Hiroyuki Takata, Mr. Takuya Itaki, and Mr. Jun'ichi Uchida of the Laboratory of Geosphere Science, Graduate School of Environmental Earth Science, Hokkaido University for their encouragement, and help in collecting references.

Deep appreciation is expressed to Mr. Yasuhumi Terasaki, the chief curator of the Imakane Board of Education, for his kind intentions during the course of this study.

## References

- Asano K., 1936a, Foraminifera from Kuromatsunai-mura, Suttu-gun, Hokkaido. Studies on the fossil Foraminifera from the Neogene of Japan. (II). Jour. Grol. Soc. Japan, 43, 419-422. (in Japanese)
- Asano K., 1936b, Foraminifera from Kuromatsunai-mura, Suttu-gun, Hokkaido. Studies on the fossil Foraminifera from the Neogene of Japan. Part 2. Jour. Geol. Soc. Japan, 43, 615-624.
- Asano K., and Nakamura M., 1937, On the distribution of the Japanese species of Cassidulina. Jour. Geol. Soc. Japan, 44, 42-49. (in Japanese)
- Asano K., 1937, Pliocene foraminifera from the Setana Beds, Hokkaido. Jour. Geol. Soc. Japan, 44, 509-512. (in Japanese)

## Foraminifera from the Kuromatsunai and Setana Formations

- Asano K., 1939, Pliocene (Upper Mizuho) Foraminifera from Japan (preliminary Report). Jour. Geol. Soc. Japan, 46, 155-168. (in Japanese with English abstract)
- Hashimoto W., Kanno S., Shinada Y., and Oshima K. , 1963, Geology of the Imagane, Kun'nui, and Yakumo districts, Oshima Peninsula, Hokkaido. Jour. Geol. Soc. Japan, 69, 228-238. (in Japanese with English abstract)
- Kato M., Katsui Y., Kitagawa Y., and Matsui M. (eds), 1990, Regional geology of Japan. Part 1. Hokkaido. Kyoritsu Shuppan, 337p.
- Masatani K., and Ohkura T., 1980MS, Neogene biostratigraphy in Oshima Peninsula, Hokkaido, Japan. In particular relation between "*Operculina-Miogypsina* Zone" and Planktonic foraminifera. Technol. Lab. JAPEX, 23, 32-52. (in Japanese with English abstract)
- Matsui M., Doi S., Takeda H., Uozumi S., Hujie T., Akiba C., Yoshimura N., Yamaguchi K., Obara T., and Odagiri T. , 1955, Report on geology and ore depositions in the north-eastern Imakane area, Shiribeshi, Hokkaido, Japan. Imakane town, 56. (in Japanese)
- Nagao T., and Sasa Y., 1933, The Cenozoic systems of southwestern Hokkaido and its geological history, part 1-2. Jour. Geol. Soc. Japan, 40 555-577; 750-775. (in Japanese)
- Nojo A., Togo, Y., Suzuki, A. and Sagayama, T., 1994, Lithostratigraphy and chronology of the Neogene Kuromatsunai Formation in the Imakane area, southwestern Hokkaido, Japan. Jour. Geol. Soc. Japan, 100, 771-786. (in Japanese with English abstract)
- Nojo A., Uchida J., Soeda Y., Itagaki K., Nakagawa M., Nikaido A., Suzuki A. and Togo Y., 1997, Some Plio-Pleistocene foraminifera from the Imakane-Kamiyakumo area of southwestern Hokkaido, Japan (Preliminary report). Bulletin of the studies in Imakane district, no. 3, 29-37. (in Japanese)
- Nojo A., Hasegawa S., Okada H., Togo Y., Suzuki A., and Matsuda T., in press, Interregional lithostratigraphy and biostratigraphy of the Pleistocene Setana Formation, southwestern Hokkaido, Japan. Geol. Soc. Japan. (in Japanese with English abstract)
- Okada T., and Mitani K. , 1981, Geology of the Imakane town. Imakane town, 77. (in Japanese)
- Shirai T., 1959, Geology in the Kuromatsunai district, with a special reference to the fossil foraminifera of the Pliocene Setana Formation—A study of the Neogene fossil foraminifera in northern Japan (1). Bull. Geol. Comm. Hokkaido, 38, 1-12. (in Japanese with English abstract)
- Shirai T., 1960, New genus and species of foraminifera from the Pliocene formation, southwestern Hokkaido. Jour. Fac. Sci. Hokkaido Univ., Ser. 4, 10, 537-543.

## Plates (Scale bar = 100μm)

### Planktonic Foraminifera (Plate 1)

1. *Globigerina bulloides* d'Orbigny, 2. *Globigerina quinqueloba* Natland,
3. *Globigerinita uvula* (Ehrenberg), 4. *Globigerinita glutinata* (Egger),
5. *Globorotalia inflata* (d'Orbigny), 6. *Globigerinoides ruber* (d'Orbigny),
- 7-9. *Neogloboquadrina incompta* (Chifelli), 10-11. *Neogloboquadrina pachyderma* (Ehrenberg)

### Benthic Foraminifera (Plate 2-12)

#### Plate 2

1. *Angulogerina kokozuraensis* (Asano), 2. *Uvigerina akitaensis* Asano,
3. *Uvigerina schencki* Asano, 4. *Uvigerina juncea* Cushman and Todd,
5. *Rectobolivina rephanus* (Parker and Jones), 6. *Buliminella elegantissima* (d'Orbigny),
7. *Bolivinita quadrilatera* (Schwager), 8. *Spiroplectammina* sp. A, 9. *Bolivina* sp. A,
10. *Bolivina decussata* Brady, 11. *Spirillina limbata* Brady,
12. *Cancris auriculus* (Fichtel and Moll)

#### Plate 3

1. *Oolina melo* d'Orbigny, 2-3. *Lagena apiopleura* Loeblich and Tappan,
4. *Oolina costata* (Williamson), 5. *Fissurina lacunata* (Burrows and Holland),
6. *Lagena striata* (d'Orbigny), 7. *Oolina striatopunctata* (Parker and Jones),
8. *Lagena acuticosta* Reuss, 9. *Fissurina baccata* (Hellon-Allen and Earland),
10. *Fissurina orbigniana* Seguenza, 11. *Fissurina* cf. *rizzae* Seguenza,
12. *Fissurina annectens* (Burrows and Holland), 13. *Fissurina* cf. *subquadrata* Parr,
14. *Fissurina marginata* (Montagu)

#### Plate 4

1. *Cibicides lobatulus* (Warker and Jacob), 2-3. *Cibicides refulgens* de Montfort,
- 4-5. *Cibicides subdepressus* (Asano), 6. *Heterolepa subhaidingerii* (Parr),
7. *Anomalinoides globulosus* (Cushman and Parr),
8. *Dyocibicides perforata* Cushman and Valentine

#### Plate 5

1. *Buccella frigida* (Cushman), 2. *Buccella makiyamai* (Chiji), 3. *Buccella tenerrima* (Brady),
4. *Patellinella hanzawai* Asano, 5. *Hanzawaia nipponica* Asano, 6. *Rosalina australis* (Parr),

Foraminifera from the Kuromatsunai and Setana Formations

7. *Rosalina bradyi* (Cushman), 8. *Guttulina yabei* Cushman and Ozawa

Plate 6

1. *Elphidium subgranulosum* Asano, 2. *Elphidium frigidum* Cushman,
- 3-5. *Elphidium subarcticum* Cushman, 6-7. *Elphidium crispum* (Linné),
8. *Polystomelina discorbinooides* Yabe and Hanzawa, 9. *Elphidium jensenii* (Cushnan),
10. *Criboelphidium yabei* (Asano), 11. *Criboelphidium oregonensis* (Cushman and Grant)

Plate 7

- 1, 8. *Astrononion hamadaense* Asano,
2. *Elphidium excavatum* (Terquem) forma *clavata* Cushman,
3. *Nonion japonicus* Asano, 4. *Melonis uchioi* Hasegawa, 5. *Pullenia apertura* Cushman,
6. *Melonis pompiliooides* (Fichtel and Moll), 7. *Lenticulina nicobarensis* (Schwager),
9. *Nonionellina labradorica* (Dawson)

Plate 8

1. *Planoglabratella?* sp. C, 2. *Planoglabratella australensis* (Hellon-Allen and Earland),
3. *Planoglabratella opercularis* (d'Orbigny), 4. *Planoglabratella subopercularis* (Asano),
5. *Glabratella mirabiris panamensis* Seiglie and Bermúdez,
6. *Planoglabratella patelliformis* (Brady), 7-8. *Glabratella* sp. A,
9. *Glabratella pulvinata* (Brady), 10. *Glabratella* sp. B

Plate 9

1. *Planoglabratella subopercularis* (Asano), 2. *Epistominella pulchella* Husezima and Maruhasi,
3. *Ammonia beccarii* (Linné), 4. *Islandidlla norcrossi* (Cushman),
5. *Islandiella japonica* (Asano and Nakamura), 6. *Islandiella yabei* (Asano and Nakamura),
7. *Islandiella helanae* Feyling-Hanssen and Buzas,
8. *Islandiella sublimbata* (Asano and Nakamura),
9. *Pseudoparella takayanagii* (Iwasa), 10. *Pseudoparella naraensis* Kuwano

Plate 10

1. *Quinqueloculina* sp.G, 2. *Pateoris hauerinoides* (Rhumbler),
3. *Quinqueloculina costata* d'Orbigny, 4. *Quinqueloculina kuromatunaiensis* Asano,
5. *Quinqueloculina agglutinata* Cuslunan, 6. *Quinqueloculina* sp.F

Plate 11

1. *Cycloforia contrita* (d'Orbigny), 2. *Quinqueloculina akneriana* d'Orbigny,


3. *Quinqueloculina* sp. C, 4. *Cribrolinoides curta* (Cushman), 5. *Quinqueloculina* sp. B,
6. *Quinqueloculina yessoensis* Asano


**Plate 12**

1. *Quinqueloculina akneriana* d'Orbigny, 2. *Quinqueloculina sawanensis* Asano,
3. *Triloculina suttuensis* Asano, 4. *Triloculina tricarinata* d'Orbigny, 5-6. *Pyrgo ezo* Asano

Foraminifera from the Kuromatsuai and Setana Formations


Plate 1





Foraminifera from the Kuromatsunai and Setana Formations


Plate 3






Foraminifera from the Kuromatsunai and Setana Formations


Plate 5






Foraminifera from the Kuromatsunai and Setana Formations


Plate 7






Foraminifera from the Kuromatsunai and Setana Formations

Plate 9





Foraminifera from the Kuromatsunai and Setana Formations

Plate 11

