



Foraminifera from the Kuromatsunai and Setana Formations in southwestern Hokkaido, Japan -Preliminary report-

メタデータ	言語: English 出版者: 公開日: 2012-10-29 キーワード (Ja): キーワード (En): 作成者: 能條, 歩 メールアドレス: 所属:
URL	https://hokkyodai.repo.nii.ac.jp/records/9395

Foraminifera from the Kuromatsunai and Setana Formations in southwestern Hokkaido, Japan -Preliminary report-

Ayumu NOJO¹⁾

Key words : foraminifera, Kuromatsunai Formation, Setana Formation, Pliocene, Pleistocene,
southwestern Hokkaido.

1. Introduction

Neogene to Quaternary deposits distributed along the Japan Sea side in Hokkaido, implying a history of development and changes of the Japan Sea, are one of most important subjects for the study to understand geological and geographical frameworks of the present Japanese Islands. In spite of these facts, there have not been enough data about the environmental changes of these areas, near the northern end of the Japan Sea areas, or around the Oshima Peninsula.

The Oshima Peninsula in southwestern Hokkaido is regarded geologically as the northern extension of the "Green Tuff" region of the Inner Zone of the Northeast Japan (e.g. Kato et al. eds., 1990). The peninsula has extensive distributions of the Neogene and Quaternary deposits which unconformably overlie pre-Tertiary sedimentary and igneous rocks. This region has been one of stratigraphical standards of the Neogene in Hokkaido since the study by Nagao and Sasa (1933).

Neogene to Pleistocene marine sediments in this region are generally classified into the following ascending-ordered formations, Kunnui, Yakumo, Kuromatsunai, and Setana (Nagao and Sasa, 1933). These formations are well exposed in middle to northern parts of the peninsula. Such stratigraphy first proposed by Nagao and Sasa (1933) has been accepted, but recent biostratigraphical data reveal that the Setana Formation should be of Pleistocene age (e.g. Nojo et

¹⁾ Department of Science Education, Iwamizawa College, Hokkaido University of Education, Iwamizawa. 068-8642 Japan.

al., in press). General relationships among these formations are as follows: from the Kunnui to Kuromatsunai Formation they contact conformably each other, but they are clinounconformably overlain by the Setana (e.g. Nagao and Sasa, 1933; Matsui et al., 1955; Hashimoto et al., 1963; Oka and Mitani, 1981; Nojo et al., 1994).

There are a few reports on microfossils in the Kuromatsunai and Setana Formations. In particular, despite abundant yieldings of foraminifera from the Setana Formation, there have not been so many reports on foraminifera (Asano, 1936a, b, 1937, 1939; Asano and Nakamura, 1937; Shirai, 1959, 1960; Masatani and Ohkura, 1980MS; Takahashi et al., 1980MS; Tsuchi and Ibaragi, 1982; Tsubakihara et al., 1989; Nojo et al., 1997), and we have only chronostratigraphical data by Nojo et al. (in press), which are related to paleoenvironmental changes on the basis of micropaleontological analyses.

The rock materials for this study were sampled from the formations in northern to middle part of the Oshima Peninsula, and were analyzed in 1994 to 1998 by the author when he was a curator of Imakane Borad of Education. Some data of these works have been reported by Nojo et al. (1997: Tab. 1). This study is preliminary report and list of foraminifera from the Kuromatsunai and Setana Formations to indicate a history of paleoenvironmental changes. The author mainly describes about some occurrences of foraminifera (Tab. 2) and shows their faunal list with plates in this report. A detailed discussion about paleoenvironmental changes is left for another opportunity.

Table 1 Foraminifera from the Kuromatsunai and Setana Formation in Imakane area (after Nojo et al., 1997)

Sample number (Kmu: Sakkurubetsu Pyro. Mem., Kms: Okusawa Sand. Mem. Ss: Setana F.)	Kmu-1	Kmu-2	Kmu-3	Kmu-4	Kmu-5	Kms-1	Ss-1	Ss-2	Sample number (Kmu: Sakkurubetsu Pyro. Mem., Kms: Okusawa Sand. Mem. Ss: Setana F.)	Kmu-1	Kmu-2	Kmu-3	Kmu-4	Kmu-5	Kms-1	Ss-1	Ss-2	
Planktonic Foraminifera									Benthic Foraminifera									
<i>Globigerina bulloides</i> d'Orbigny	+	+	+	+				+	<i>Fissurina</i> cf. <i>rizzae</i> Seguenza				+					
<i>Globigerina quinqueloba</i> Nalband			+					+	<i>Fissurina</i> cf. <i>subquadrata</i> Parr		+							
<i>Globigerina</i> sp.								+	<i>Fissurina</i> sp.			+						
<i>Globigerinita glutinata</i> (Egger)		+						+	<i>Glabratalia</i> spp.		+	+					+	
<i>Globigerinita uvula</i> (Ehrenberg)								+	<i>Globocassidulina bisecta</i> Nomum		+	+	+					
<i>Neogloboquadrina incompta</i> (Giffeli)								+	<i>Globocassidulina jamesoni</i> (McCulloch)				+					
<i>Neogloboquadrina pachyderma</i> (Ehrenberg) (dextral)	+	+	+	+				+	<i>Globocassidulina</i> spp.			+					+	
<i>Neogloboquadrina pachyderma</i> (Ehrenberg) (sinistral)	+	+	+	+				+	<i>Gutulina yabei</i> Cushman & Ozawa								+	
Benthic Foraminifera									<i>Gutulina</i> sp.									
<i>Ammonia</i> sp.								+	<i>Gutulina</i> spp.		+	+					+	+
<i>Angulogerina kokuzuraensis</i> Asano				+					<i>Hanzawaia nipponica</i> Asano									
<i>Angulogerina</i> sp.			+						<i>Heterolepa praecincta</i> (Karrer)			+						
<i>Asteronion hamadaense</i> Asano				+					<i>Heterolepa subhaidingerii</i> (Parr)				+					
<i>Bolivina decussata</i> Brady					+				<i>Islandiella japonica</i> (Asano & Nakamura)		+		+				+	+
<i>Bolivina</i> spp.				+	+				<i>Islandiella norcrossi</i> (Cushman)			+						
<i>Bolivinita quadrilatera</i> (Schwager)		+							<i>Islandiella setanaensis</i> (Asano & Nakamura)		+		+				+	+
<i>Buccella frigida</i> (Cushman)				+	+				<i>Islandiella subimbata</i> (Asano & Nakamura)		+	+	+				+	+
<i>Buccella tenerima</i> (Bandy)				+					<i>Islandiella yabei</i> (Asano & Nakamura)			+	+	+			+	+
<i>Buccella</i> sp.				+					<i>Islandiella</i> spp.		+	+					+	+
<i>Bulimina</i> sp.								+	<i>Lagena acutirostris</i> Reuss		+							
<i>Cibicides lobatulus</i> (Walker & Jacob)				+				+	<i>Lagena apioplura</i> Loeblich & Tappan				+					
<i>Cibicides refulgens</i> de Montfort				+				+	<i>Lagena striata</i> (d'Orbigny)		+							
<i>Cibicides subdepressus</i> (Asano)				+				+	<i>Lagena</i> spp.				+					
<i>Cibicides</i> spp.				+	+			+	<i>Lenticulina</i> sp.		+							
<i>Criboelphidium yabei</i> (Asano)				+				+	<i>Melonis pompilioides</i> (Fichtel & Moll)		+							
<i>Criboelphidium</i> sp.								+	<i>Melonis uchii</i> Hasegawa		+							
<i>Dyocibicides perforata</i> Cushman & Valentine		+						+	<i>Nonion</i> sp.			+						
<i>Elphidium crispum</i> (Linné)				+				+	<i>Nonionella labradorica</i> (Dawson)					+	+			
<i>Elphidium excavatum</i> (Terquem) forma <i>clavata</i> (Cushman)				+	+			+	<i>Oolina striatopunctata</i> (Parker & Jones)					+	+			
<i>Elphidium jenseni</i> (Cushman)				+				+	<i>Ordosiralis umbonatus</i> (Reuss)		+							
<i>Elphidium subarcticum</i> Cushman				+				+	<i>Planoglobatella subopercularis</i> (Asano)		+							
<i>Elphidium</i> spp.				+				+	<i>Polysiomellina discorbinoidea</i> Yabe & Hanzawa			+						
<i>Epistominella pulchella</i> Husezima & Maruhasi				+				+	<i>Pseudoparrella takayanagii</i> (Iwasa)				+					
<i>Epistominella</i> spp.				+				+	<i>Pullenia apertura</i> Cushman				+					
<i>Eponides</i> ? sp.				+	+			+	<i>Pyrgo</i> <i>ezo</i> Asano					+				
<i>Fissurina annectens</i> (Burrows & Holland)				+				+	<i>Silostomella</i> sp.		+							
<i>Fissurina baccata</i> (Heron-Allen & Earland)		+						+	<i>Uvigerina aktaiensis</i> Asano					+				
<i>Fissurina marginata</i> (Montagu)		+	+	+	+			+	<i>Uvigerina juncea</i> Cushman & Todd		+							
<i>Fissurina orbignyana</i> Seguenza		+	+	+	+			+	<i>Uvigerina schencki</i> Asano				+					
									<i>Uvigerina</i> spp.									

Foraminifera from the Kuromatsunai and Setana Formations

Table 2 Foraminifera from the Kuromatsunai and Setana Formations

Sampling horizon	Upper Kuroma Tsu-nai F.	Lower Setana F.	Upper Setana F.	Sampling horizon	Upper Kuroma Tsu-nai F.	Lower Setana F.	Upper Setana F.
Planktonic Foraminifera				Benthic Foraminifera			
<i>Globigerina bulloides</i> d'Orbigny	A	A	A	<i>Discorbinella bertheloti</i> (d'Orbigny)	-	-	R
<i>Globigerina quinqueloba</i> Natland	R	C	F	<i>Discorbinella</i> spp.	R	-	VR
<i>Globigerinita glutinata</i> (Egger)	F	F	F	<i>Eilohedra</i> spp.	R	-	-
<i>Globigerinita uvula</i> (Ehrenberg)	R	VR	F	<i>Elphidiella arctica</i> (Parker & Jones)	C	-	-
<i>Globigerinoides ruber</i> (d'Orbigny)	-	-	R	<i>Elphidium aculeatum</i> (d'Orbigny)	-	-	VR
<i>Globorotalia inflata</i> (d'Orbigny)	C	C	VR	<i>Elphidium advena</i> (Cushman)	-	-	VR
<i>Grobigerina</i> spp.	-	-	R	<i>Elphidium bartletti</i> Cushman	R	F	R
<i>Neogloboquadrina</i> cf. <i>asanoi</i> (Maiya, Saito, and Sato)	VR	-	-	<i>Elphidium crispum</i> (Linné)	F	F	F
<i>Neogloboquadrina dutertrei</i> (d'Orbigny)	VR	-	-	<i>Elphidium excavatum</i> (Terquem) forma <i>clavata</i> Cushman	-	R	F
<i>Neogloboquadrina incompta</i> (Cifelli)	C	A	A	<i>Elphidium frigidum</i> Cushman	R	-	R
<i>Neogloboquadrina pachyderma</i> (D) (Ehrenberg)	F	A	C	<i>Elphidium hanzawai</i> Asano	-	VR	-
<i>Neogloboquadrina pachyderma</i> (S) (Ehrenberg)	F	C	C	<i>Elphidium incertum</i> (Williamson)	-	-	VR
<i>Neogloboquadrina</i> sp.	-	VR	R	<i>Elphidium janseni</i> (Cushman)	VR	-	R
Gen. et sp. indet.	VR	R	VR	<i>Elphidium macellum</i> (Fichtel & Moll)	VR	-	-
TOTAL COUNTED	296	308	5035	<i>Elphidium subarcticum</i> Cushman	-	F	C
Benthic Foraminifera				<i>Elphidium subgranulosum</i> Asano	-	-	F
<i>Alabamina japonica</i> (Asano)	VR	-	-	<i>Elphidium subincertum</i> Asano	-	R	-
<i>Ammonia beccarii</i> (Linné)	-	VR	-	<i>Elphidium translucens</i> Natland	-	-	R
<i>Ammonia ketenziensis</i> (Ishizaki)	-	VR	-	<i>Elphidium</i> spp.	F	VR	R
<i>Ammonia inflata</i> (Seguenza)	-	R	-	<i>Epistominella pulchella</i> Husezima & Maruhasi	R	VR	R
<i>Ammonia takanabensis</i> (Ishizaki)	-	VR	-	<i>Epistominella</i> spp.	VR	-	R
<i>Ammonia</i> spp.	-	VR	R	<i>Eponides?</i> sp.	R	-	R
<i>Angulogerina hughesi</i> (Galloway and Wissler)	R	VR	R	<i>Fissurina annectens</i> (Burrows & Holland)	-	-	R
<i>Angulogerina kokozuraensis</i> Asano	R	-	R	<i>Fissurina</i> cf. <i>annectens</i> (Burrows & Holland)	VR	-	-
<i>Anomalina rostrata</i> (Brady)	-	-	VR	<i>Fissurina lacunata</i> (Burrows & Holland)	-	-	R
<i>Anomalina</i> spp.	-	-	R	<i>Fissurina lucida</i> (Williamson)	-	-	R
<i>Anomalinoides globulosus</i> (Chapman & Parr)	VR	-	R	<i>Fissurina marginata</i> (Walker & Boys)	R	R	F
<i>Astacolus hyalaculus</i> Loeblich and Tappan	VR	-	VR	<i>Fissurina obscurocostata</i> Galloway & Wissler	VR	-	R
<i>Astrononion aomoriense</i> Asano	-	VR	-	<i>Fissurina orbignyana</i> Seguenza	R	-	VR
<i>Astrononion hamadaense</i> Asano	-	F	R	<i>Fissurina</i> cf. <i>rizzae</i> Seguenza	-	-	R
<i>Astrononion</i> spp.	R	-	VR	<i>Fissurina semimarginata</i> (Reuss)	VR	R	-
<i>Bolivina decussata</i> Brady	F	-	R	<i>Fissurina</i> cf. <i>subquadrata</i> Parr	-	-	R
<i>Bolivina</i> sp. A	VR	R	R	<i>Fissurina</i> spp.	R	VR	R
<i>Bolivina</i> spp.	R	-	R	<i>Gavelinopsis praegeri</i> (Heron-Allen & Earland)	-	-	R
<i>Brizalina alata</i> (Seguenza)	-	-	VR	<i>Gavelinopsis</i> spp.	R	-	R
<i>Brizalina</i> spp.	-	-	R	<i>Glabratella</i> cf. <i>aurantista</i> Seiglie & Bermúdez	-	-	R
<i>Buccella frigida</i> (Cushman)	VR	R	R	<i>Glabratella murabilis panamensis</i> Seiglie & Bermúdez	-	-	R
<i>Buccella makiyamai</i> (Chiji)	VR	R	C	<i>Glabratella pulvinata</i> (Brady)	-	-	R
<i>Buccella nipponica</i> (Husezima & Maruhasi)	-	VR	R	<i>Glabratella</i> sp. A	-	-	R
<i>Buccella tanaii</i> (Uchio)	-	-	R	<i>Glabratella</i> sp. B	-	-	R
<i>Buccella tenerima</i> (Bandy)	R	-	C	<i>Glabratella</i> spp.	VR	-	R
<i>Buccella</i> spp.	VR	VR	R	<i>Glandulina ovula</i> d'Orbigny	-	-	-
<i>Bulimina</i> cf. <i>imamurai</i> Tai	R	-	-	<i>Globocassidulina bisecta</i> Nomura	F	R	R
<i>Buliminella elegantissima</i> (d'Orbigny)	-	-	F	<i>Globocassidulina canalisurata</i> Eade	-	R	-
<i>Cancris aurculus</i> (Fichtel & Moll)	-	-	VR	<i>Globocassidulina neobrocha</i>	-	-	R
<i>Chrysalidinella dimorpha</i> (Brady)	-	-	VR	<i>Globocassidulina</i> spp.	F	R	F
<i>Cibicides</i> cf. <i>kamadae</i> Asano	-	-	VR	<i>Guttulina</i> spp.	VR	-	R
<i>Cibicides lobatulus</i> (Walker & Jacob)	A	A	F	<i>Gyroidina</i> spp.	-	-	R
<i>Cibicides refulgens</i> de Montfort	C	C	C	<i>Hanzawaia nipponica</i> Asano	R	VR	R
<i>Cibicides subdepressus</i> (Asano)	R	F	R	<i>Haynesina</i> sp.	-	-	R
<i>Cibicides tani</i> Iwasa and Kikuchi	F	R	R	<i>Heronallenia</i> sp.	-	-	VR
<i>Cibicides</i> spp.	R	R	R	<i>Heterolepa subhaidingerii</i> (Parr)	-	-	R
<i>Cibicoides pachydermus</i> (Rzehak)	R	-	VR	<i>Heterolepa</i> sp.	VR	-	-
<i>Cornuspira involvens</i> (Reuss)	-	-	R	<i>Islandiella helenae</i> Feyling-Hassen & Buzas	-	-	R
<i>Criboelphidium oregonense</i> (Cushman & Grant)	-	-	VR	<i>Islandiella islandica</i> (Nørbang)	F	-	-
<i>Criboelphidium yabei</i> (Asano)	-	-	R	<i>Islandiella japonica</i> (Asano & Nakamura)	F	C	R
<i>Cribrulinoides curta</i> (Cushman)	-	-	VR	<i>Islandiella norcrossi</i> (Cushman)	-	-	R
<i>Cycloforia contorta</i> (d'Orbigny)	-	VR	R	<i>Islandiella sublimbata</i> (Asano & Nakamura)	A	A	F
<i>Cystammina pauciloculata</i> (Brady)	VR	-	-	<i>Islandiella yabei</i> (Asano & Nakamura)	F	F	-
<i>Discorbinella araucana</i> (d'Orbigny)	R	VR	F	<i>Islandiella</i> spp.	-	R	R

Ayumu NOJO

Table 2 Foraminifera from the Kuromatsunai and Setana Foraminifera (continued)

Sampling horizon	Upper Kuromatsunai F.	Lower Setana F.	Upper Setana F.	Sampling horizon	Upper Kuromatsunai F.	Lower Setana F.	Upper Setana F.
Benthic Foraminifera				Benthic Foraminifera			
Karrerella baccata japonica Asano	-	VR	-	Quinqueloculina akneriana d'Orbigny	R	R	R
Karrerella spp.	R	-	R	Quinqueloculina costata d'Orbigny	-	-	R
Lagena acuticosta (Reuss)	-	-	VR	Quinqueloculina elongata Natland	-	-	R
Lagena apiopleura (Loeblich & Tappan)	VR	VR	R	Quinqueloculina kuromatsunaiensis Asano	-	-	R
Lagena sulcata spicata Cushman & McCulloch	-	-	R	Quinqueloculina sawanensis Asano	-	-	R
Lagena sp.	-	-	VR	Quinqueloculina seminulum (Linné)	-	VR	R
Lenticulina nikobarensis (Schwager)	-	-	VR	Quinqueloculina vulgaris d'Orbigny	-	R	R
Lenticulina sp.	-	-	VR	Quinqueloculina cf. vulgaris d'Orbigny	-	-	VR
Melonis pacificus (Cushman)	VR	-	VR	Quinqueloculina yezoensis Asano	-	-	R
Melonis uchioi Hasegawa	-	-	VR	Quinqueloculina sp. B	-	-	R
Melonis spp.	VR	-	-	Quinqueloculina sp. C	-	-	R
Miliolinella circularis (Bornemann)	-	-	R	Quinqueloculina sp. F	-	-	R
Miliolinella spp.	-	-	R	Quinqueloculina sp. G	-	-	R
Neoconorbina stachi (Asano)	-	VR	-	Quinqueloculina sp. H	-	-	R
Neoeponides procerus (Brady)	-	R	R	Quinqueloculina spp.	R	R	F
Neoeponides spp.	VR	-	-	Rectobolivina rephanus (Parker & Jones)	-	-	VR
Nodosarina? sp.	-	-	VR	Rosalina australis (Parr)	VR	VR	R
Nonion japonicus Asano	-	-	R	Rosalina bradyi (Cushman)	VR	VR	F
Nonion spp.	R	VR	R	Rosalina isabelleana d'Orbigny	-	-	R
Nonionella stella Cushman & Moyer	-	-	R	Rosalina vilardeboana d'Orbigny	-	-	R
Nonionoides grateloupi (d'Orbigny)	-	-	VR	Rosalina spp.	R	VR	R
Oolina carteri Albani and Yassini	VR	-	-	Sigmohauerina sp.	-	-	VR
Oolina costata (Williamson)	-	-	VR	Sigmoidella pacifica Cushman & Ozawa	-	-	VR
Oolina melo d'Orbigny	-	VR	R	Sigmoidella spp.	VR	-	-
Oolina striatopunctata (Parker & Jones)	-	-	VR	Sigmoilina sigmoidea compressa Cushman	-	R	-
Oolina sp.	-	-	VR	Sigmoilina sp.	-	-	VR
Oridosalis umbonatus (Reuss)	-	-	VR	Sigmomorphina spp.	-	VR	-
Oridosalis cf. umbonatus (Reuss)	VR	-	-	Sigmomorphina semitecta terquemiana (Fornasini)	-	-	R
Osangularinella umbonifera (Cushman)	VR	-	-	Sigmomorphina trilocularis (Bagg)	VR	-	-
Paracassidulina sulcata Belford	-	-	R	Sigmovirgulina sp.	-	-	VR
Pararotalia nipponica (Asano)	-	-	R	Sphaeroidina japonica Brady	VR	-	-
Patellina corrugata Williamson	-	VR	R	Sphaeroidina spp.	-	-	-
Patellinella hanzawai Asano	-	-	R	Spirillina limbata Brady	-	-	VR
Pateoris hauerinoides (Rhmblor)	-	-	F	Spirillina vivipara Ehrenberg	-	-	R
Planoglabratella australensis (Heron-Allen & Earland)	-	-	R	Spirillina sp.	-	-	VR
Planoglabratella opercularis (d'Orbigny)	-	-	R	Spiroloculina hadai Thalmann	-	-	R
Planoglabratella pateriformis (Brady)	-	-	R	Spiroloculina sp.	-	-	VR
Planoglabratella subopercularis (Asano)	R	R	F	Spiroplectammina sp. A	-	-	R
Planoglabratella? sp. C	-	-	VR	Spiroplectammina sp.	-	-	VR
Polystomellina discorbinoidea Yabe & Hanzawa	-	VR	R	Spiroplectinella wrightii (Silvestri)	-	-	R
Poroepionides cribroreponodus Asano & Uchio	-	-	R	Textularia conica d'Orbigny	-	-	VR
Pseudononion japonicum Asano	-	VR	R	Textularia spp.	-	-	R
Pseudononion spp.	-	VR	VR	Triloculina rotunda d'Orbigny	-	VR	-
Pseudoparrella naraensis Kuwano	VR	-	F	Triloculina suttuensis Asano	-	-	R
Pseudoparrella takayanagii (Iwasa)	VR	R	F	Triloculina tricarinata d'Orbigny	-	-	VR
Pseudoparrella spp.	-	-	R	Triloculina sp. A	-	-	VR
Pseudopolymorphina sp.	-	-	VR	Uvigerina juncea Cushman	-	VR	-
Pullenia apertura Cushman	VR	-	-	Uvigerina spp.	VR	-	-
Pyrgo ezo Asano	VR	-	-	Valvulinera hamanakoensis (Ishiwada)	-	-	VR
Pyrgo fornasinii Chapman & Parr	-	R	-	Valvulinera japonica Asano	-	VR	-
Quinqueloculina agglutinata Cushman	-	-	R	Gen. et sp. indet.	R	F	F
Total counted					869	780	4652

A: 80.5-13.5%, C: 13.5-4.5%, F: 4.5-1.5%, R: <1.5%, VR: two or less occurrences, -: no occurrence

2. Faunal Reference List

Foraminiferal species from the Kuromatsunai and Setana Formations are alphabetically listed below, under the categories of planktons and benthos. The selected species are illustrated with scanning electron micrographs in Plates 1 to 12. The original references are given for species. Additional references are selected from those with helpful remarks and illustrations about the species from the areas studied in this report. All specimens are catalogued and deposited in Department of Science Education, Iwamizawa College, Hokkaido University of Education.

Planktonic Foraminifera

Globigerina bulloides d'Orbigny

Globigerina bulloides d'Orbigny, 1826, Ann. Sci. Nat., Paris, ser. 1, v. 7, p. 277, Modéle nos. 17, 76; Brady, 1884, Voy. Challenger, Rep., Zool., v. 9, p. 593, pl. 79, figs. 7a-c.

Globigerina quinqueloba Natland

Globigerina quinqueloba Natland, 1938, Bull. Scripps Inst. Oceanogr., Tech. Ser., v. 4, no. 5, p. 149, pl. 6, figs. 7a-c.

Globigerinita glutinata (Egger)

Globigerina glutinata Egger, 1893, Abhandl. K. Bayer. Akad. Wiss. Munchen, CLII, v. 18, p. 371, pl. 13, figs. 19-21.

Globigerinita glutinata (Egger), Parker, 1962, Micropal. v. 8, no. 2, p. 246, pl. 9, figs. 1-16.

Globigerinita uvula (Ehrenberg)

Pyloedixia uvula Ehrenberg, 1861, K. Preuss., Akad. Wiss. Berlin, Monatsber., p. 276, 277, 308.

Globigerinita uvula (Ehrenberg), Parker, 1962, Micropal., v. 8, no. 2, p. 252, pl. 8, figs. 14-26.

Globigerinoides ruber (d'Orbigny)

Globigerina rubra d'Orbigny, 1839, in Sagra, R. de la, Hist. Ohys. Pol. Nat. Cuba, Foraminiferes, p. 82, pl. 4, figs. 12-14.

Globigerinoides rubra (d'Orbigny) (sic.), Cushman, 1927, Cushman Lab. Foram. Res., Contr., v. 3, pt. 1, p. 87, pl. 19, figs. 6a-c.

Globorotalia inflata (d'Orbigny)

Globigerina inflata d'Orbigny, 1839, in Barker-Webb and Berthelot, Hist. Nat. Canaries, v. 2, pt. 2, Zool., p. 134, pl. 2, figs. 7-9.

Globorotalia inflata (d'Orbigny), sensu strict, Maiya Saito and Sato, 1976, in Takayanagi and Saito eds. Progress in Micropaleontology, Micropal. Press, p. 408, pl. 2, figs. 5-7.

Neogloboquadrina cf. *asanoi* (Maiya, Saito, and Sato)

cf. *Globoquadrina asanoi* Maiya, Saito, and Saito, 1976, Progress in Micropaleontology. p.

409, pl. 3, figs. 1a-c, 2a-c, 3.

cf. *Neogloboquadrina asanoi* (Maiya, Saito, and Sato), Thompson, 1980, Init. Rep. DSDP, v. 56-57, pl. 3, figs. 10-12.

Neogloboquadrina dutertrei (d'Orbigny)

Globigerina dutertrei d'Orbigny, 1839, Foraminiferés, p. 84, pl. 4, figs. 19-21.

Neogloboquadrina dutertrei (d'Orbigny), Rögl and Bolli, 1973, Init. Rep. DSDP, v. 27, 743-767, pl. 9, figs. 1-3, 7-10; pl. 17, figs. 1-6.

Neogloboquadrina incompta (Cifelli)

Globigerina incompta Cifelli, 1961, Cushman Found. Foram. Res., Contr., v. 12, pt. 3, p. 84, pl. 4, figs 1-7.

Neogloboquadrina incompta (Cifelli), Rogl and Bolli, 1973, Init. Repts. DSDP, v. 15, p. 571, pl. 10?

Neogloboquadrina pachyderma (Ehrenberg)

Aristerospira pachyderma Ehrenberg, 1861, K. Preuss. Akad. Wiss, Berlin, Monatsber. p. 276, 277, 303.

Globigerina pachyderma (Ehrenberg), Brady, 1884, Voy. Challenger, Rep., Zool., v. 9, p. 600, pl. 114, figs. 19, 20.

Neogroboquadrina pachyderma (Ehrenberg), Rogl and Bolli, 1973, Init. Repts. DSDP, v. 15, p. 571, pl. 11, figs. 2-6; pl. 16, fig. 12.

Benthic Foraminifera

Alabamina japonica (Asano)

Pseudoparrella japonica Asano, 1949, Jour. Paleont., 23 (4), p430, fig. 2, Nos. 2-4.

Alabamina japonica (Asano), Takayanagi and Hasegawa, 1987, Checklist and bibliography of post-Paleozoic foraminifera established by Japanese workers, 1890-1986, p. 39.

Ammonia beccarii (Linné)

Nautilus beccarii Linne, 1758, Syst. Nat., se. 10, p. 710.

Ammonia beccarii (Linne), Fizzell and Keen, 1949, Jour. Pal., v. 23, no. 1, p. 106.

Ammonia ketienziensis (Ishizaki)

Stebulus ketienziensis Ishizaki, 1948, Acta Geol. Taiwan, 2 (1), 59, pl. 1, figs. 2a-c.

Ammonia ketienziensis (Ishizaki), Huang, 1964, Micropaleont. v. 10, no. 1, p. 53, pl. 1, figs. 13a-c.

Angulogerina hughesi (Galloway and Wissler)

Uvigerina hughesi Galloway and Wissler, 1927, Jour. Pal., v. 1, p.76, pl. 12, figs. 5a, b.

Angulogerina hughesi (Galloway and Wissler), Asano, 1950, Pacific Sci. v. 4, no. 2, p. 19, figs. 91-93.

Angulogerina kokozuraensis Asano

Angulogerina kokozuraensis Asano, 1949, Jour. Pal., v. 23 no. 4, p. 428, text-fig. 1, nos. 50-53; 1950, Illust. Cat. Japan. Tert. Small. Foram., pt. 2, p. 19, figs. 94-96.

Trifarina kokozuraensis (Asano), Matoba, 1967, Tohoku Univ., Sci. Rep., 2nd ser. (Geol.) ,

Foraminifera from the Kuromatsunai and Setana Formations

v.38, no.2, p. 257, pl. 26, fig.3.

Anomalinella rostrata (Brady)

Truncatulina rostrata Brady, 1881, Quart. Jour. Micr. Sci., n. s., v. 21, p. 65.

Anomalinella rostrata (Brady), Cushman, 1927, Cont. Cushman Labo. Foram. Res., v. 3, p. 93.

Anomalinoides globulosus (Chapman and Parr)

Anomalina grosserugosa Brady, 1884, Challenger Foram., p.675, pl. 94, figs. 4, 5.

Anomalina globulosa Chapman and Parr, 1937, Austr. Antarct. Exp. 1911-14, Sci. Rep., ser. C, 1 (2), p.117.

Astacolus hyalacrulus Loeblich and Tappan

Astacolus hyalacrulus Loeblich and Tappan, 1953, Smithsonian Misc. Coll. v. 121, no. 7, p.52, pl. 9, figs. 1-4.

Astrononion aomoriense Asano

Astrononion aomoriense Asano, 1950, Illust. Cat. Japan. Tert. Small. Foram., pt.1, p. 5, figs. 25, 26.

Astrononion hamadaense Asano

Astrononion hamadaense Asano, 1950, Illust. Cat. Japan. Tert. Small. Foram., pt.1, p.6, figs. 29-31.

Bolivina decussata Brady

Bolivina decussata Brady, 1881, Quart. Jour. Micr. Sci., n.s., v. 21, p. 28; 1884, Voy. Challenger, Rep., Zool., v. 9, p. 423, pl. 53, figs. 12-13.

Bolivina sp.A

This form is identified to *Bolivina* sp. A of Hasegawa (1979), and is similar to *Bolivina decussata* Brady in its outline, but differs from it in less developed, and rounded lobes on the wall surface.

Bolivinita quadrilatera (Schwager)

Textularia quadrilatera Schwager, 1866. Novara Exp. Geol. Theil., 2, p. 253, pl. 7, fig. 10.

Bolivina quadrilatera (Schwager), Asano, 1938, Jour. Geol. Soc. Japan, 45, p. 607, pl.16, fig. 10.

Brizalina alata (Seguenza)

Vulvulina alata Seguenza, 1862, Atti Accad. Gioenia Sci. Nat., 2 (18), p. 115, pl. 2, figs.5, 5a.

Bolivina alata (Seruenza), Cushman, 1937, Cushman Lab. Foram., Res. Spac. Publ., 9, p. 106, pl.13, figs. 3-11.

Brizalina alata (Seguenza), Belford, 1966, Bureau of Mineral Resources, Geol. and Geophysics, Bull. 79.

Buccella frigida (Cushman)

Pulvinulina frigida Cushman, 1922, Canada Biol. Contr., no. 9 (1921), p. 12.

Buccella frigida (Cushman), Andersen, 1952, Washington Acad. Sci., Jour., v. 42, no. 5, p. 144, figs. 4-6.

Buccella kuromatsunaiensis Shirai, 1960, Jour. Fac. Sci. Hokkaido Univ., Ser. IV, vol. X, 537-543, pl. 2, figs. 2a-c.

Buccella makiyamai Chiji

- Eponides schreibersii* Morishima and Chiji (not of Reuss), 1952, Mem. Coll. Sci., Univ. Kyoto, ser. B, v. 20, no. 2, pl. 2 (XIII), figs. 6a-c.
- Buccella makiyamae* (sic.) Chiji, 1961, Prof. J. Makiyama, Mem. Vol., Kyoto, p. 234, text-figs. 2a-c, pl. 1, figs. 13-14.
- Buccella nipponica* (Husezima and Maruhasi)
- Discorbis nipponica* Husezima and Maruhasi, 1944, Jour. Shigenkagaku Kenkyusho (Res. Inst. Nat. Resour., Japan), v. 1, no. 3, p. 397, pl. 34, figs. 9a-c.
- Buccella nipponica* (Husezima and Maruhasi), Hasegawa, 1979, Tohoku Univ., Sci. Rep., 2nd ser. (Geol.), v. 49, no. 2, p. 144, pl. 7, figs. 4a-d.
- Buccella tanaii* (Uchio)
- Eponides tanaii* Uchio, 1951, Jour. Geol. Soc. Japan, 57, p. 376, pl. 5, fig. 8.
- Buccella tanaii* (Uchio), Takayanagi and Hasegawa, 1987, Checklist and bibliography of post-Paleozoic foraminifera established by Japanese workers, 1890-1986, p. 19.
- Buccella tenerrima* (Bandy)
- Rotalia tenerrima* Bandy, 1950, Paleont., vol. 24, Tulsa, Okula., p. 278, pl. 42, fig. 3.
- Eponides frigidus* (Cushman), Cushman and Todd, 1947, (not *Pulvinulina frigida* Cushman, 1922), Cushman Lab. Foram. Res. Spec. Publ. v. 23, p. 71, pl. 8, fig. 7.
- Buccella inusitata* Andersen, 1952, Washington Acad. Sci., Jour., v. 42, no. 5, p. 148, figs. 10-11.
- Buccella tenerrima* (Bandy), Feyling-Hanssen, 1976, Maritime Sediments Spec. Publ. 1, p. 353, pl. 1, fig. 8-10; pl. 2, fig. 10-12.
- Bulimina* cf. *imamurai* Tai
- cf. *Bulimina imamurai* Tai, 1959, Hiroshima Univ., Jour. Sci. ser. C, v. 2, no. 4, p. 387, pl. 40, fig. 3.
- Buliminella elegantissima* (d'Orbigny)
- Bulimina elegantissima* d'Orbigny, 1839, Voy. Amer. Mérid., Foraminifères, v. 5, pt. 5, p. 51, pl. 7, figs. 13, 14.
- Buliminella elegantissima* (d'Orbigny), Cushman, 1919, U.S. Nat. Mus., Proc., v. 56, p. 606.
- Cancris auriculus* (Fichtel and Moll)
- Nautilus auriculus* Fichtel and Moll, 1789, Testacea microscopica, p. 105, pl. 20, figs. a-f; 1803 var. a, p. 108, pl. 20, figs. a-c; var. b, p. 110, pl. 20, figs. d-f.
- Cancris auricula* (Fichtel and Moll) (sic.), Cushman, 1927, Bull. Scripps Inst. Oceanogr., Tech. Ser., v. 1, no. 10, p. 164, pl. 5, fig. 10.
- Chrysalidinella dimorpha* (Brady)
- Chrysalidina dimorpha* Brady, 1881, Quart. Jour. Micr. Sci., n. s., v. 21, p. 24; 1884, Voy. Challenger, Rep., Zool., v. 9, p. 388, pl. 46, figs. 20-21.
- Chrysalidinella dimorpha* (Brady), Schubert, 1908, Neues Jb. Min. Geol. Pa., 1., Beil.-Bd. 25, p. 242.
- Cibicides* cf. *kamadai* Asano
- cf. *Cibicides kamadai* Asano, 1951, Illust. Cat. Japan. Tert. Small. Foram., pt. 13, p. 17, figs. 33-35.
- Cibicides lobatulus* (Walker and Jacob)
- Nautilus lobatulus* Walker and Jacob, 1789, Adams Essays, p. 642, pl. 14, fig. 36.

Foraminifera from the Kuromatsunai and Setana Formations

- Cibicides lobatula* (Walker and Jacob) (sic.), Cushman, 1931, U.S. Nat. Mus., Bull. 104, pt. 8, p. 118, pl. 21, figs. 3a-c.
- Cibicides refulgens* de Montfort
Cibicides refulgens de Montfort, 1808, Conch. Syst., v. 1, p. 122.
Truncatulina refulgens (Montfort), Brady, 1884, Voy. Challenger, Rep., Zool., v. 9, p. 659, pl. 92, figs. 7-9.
- Cibicides subdepressus* (Asano)
Planulina subdepressa Asano (sic.), 1951, Illust. Cat. Japan. Tert. Small. Foram., pt. 13, p. 15, figs. 16-18.
Cibicides subdepressus (Asano), Takayanagi and Hasegawa, 1987, Checklist and bibliography of post-Paleozoic foraminifera established by Japanese workers, 1890-1986, p. 37.
- Cibicides tani* Iwasa and Kikuchi
Cibicides tani Iwasa and Kikuchi, 1954, Paleont. Soc. Japan, Trans. Porc., N. S., no. 16, p. 193, text-figs. 8a-c.
- Cibicidoides pachydermus* (Rzehak)
Truncatulina pachyderma Rzehak, 1886, Die Foraminiferenfauna der Neogenformation der Umgeburg von Mähr.-Osturf. Naturf. Ver. Brünn, Verh., Bruenn [Brno, Czechoslovakia], vol. 24 (1885), p. 87, pl. 1, fig. 5a-c.
Truncatulina pseudoungeriana Cushman, 1922, U.S. Geol. Surv. Prof. Pap., 129 E, p.97, pl. 20, fig. 9.
Cibicidoides pachyderma (sic.), van Morkhoven, 1986, Cenozoic Cosmopolitan Deep-Water Benthic Foraminifera, p. 68-71, pl. 22.
- Cornuspira involvens* (Reuss)
Operculina involvens Reuss, 1849, Denkschr. Akad. Wiss., Wien, vol. 1, p. 370, pl. 45, fig. 20.
Cornuspira involvens (Reuss), Barker, 1960, SEPM, Spec. Publ., 9, pl. 11, figs. 1-3.
- Criboelphidium oregonensis* (Cushman and Grant)
Elphidium oregoense Cushman and Grant (sic.), 1927, San Diego Soc. Nat. Hist., Trans., V, no.6, 79, viii, 3.
- Criboelphidium yabei* (Asano)
Elphidium yabei Asano, 1938, Geol. Soc. Japan, Jour., v. 45, no. 538, p. 589, pl. 14, figs. 9-10.
Criboelphidium yabei (Asano), Asano, 1950, Illust. Cat. Japan. Tert. Small. Foram., pt. 1, p. 11, figs. 64-65.
- Cribrulinoides curta* (Cushman)
Quinqueloculina disparilis d'Orbigny var. *curta* Cushman, 1917, U.S. Nat. Mus. Bull. v. 71, no. 6, p.49, pl. 14, fig. 2.
Cribrulinoides curta (Cushman), Cushman and Leroy, 1939, Contr. Cushman Lab. Foram. Res., v. 15, no. 1, p. 17, pl. 3, figs. 1-11; pl. 4, figs. 1-13.
- Cycloforia contrta* d'Orbigny
Cycloforia contrta d'Orbigny, 1846, Foram. Foss. Viene, p. 298, pl. 20, figs. 4-6.
- Cystamina pauciloculata* (Brady)

- Trochammina pauciloculata* Brady, 1879, Quaterly Jour. Microscopical Science, new ser. 19, p. 58, pl. 5, figs. 13, 14.
- Ampchilostoma pauciloculata* (Brady), Eimer and Fickert, 1899, Zeitschrift Wissenschaftliche Zoologie 65, p. 692.
- Cystamina pauciloculata* (Brady), Galloway, 1933, A Manual of Foraminifera. p. 186, pl. 16, fig. 12.
- Discorbinella araucana* (d'Orbigny)
Rosanila araucana d'Orbigny, 1839,
Discorbinella araucana (d'Orbigny), Jones, 1994, Challenger Foraminifera, p. 93, pl. 86, figs. 10, 11.
- Discorbinella bertheloti* (d'Orbigny)
Rosalina bertheloti d'Orbigny, 1839, in Barker-Webb and Berthelot, Hist. nat. Canaries, v. 2, pt. 2, Zool., p. 135, pl. 1, figs. 28-30
Discorbinella bertheloti (d'Orbigny), Loeblich and Tappan, 1964, Treatise on Invertebrate Pal., Moore R.C., ed., pt. C, Protista, 2, v. 2, p. C575, figs. 453.
- Dyocibicides perforata* Cushman and Valentine
Dyocibicides perforata Cushman and Valentine, 1930, Contr. Stanford Geol. Dept., v. 1, no. 1, p. 31, pl. 10, figs. 3a-c.
- Elphidiella arctica* (Parker and Jones)
Polystomella arctica Parker and Jones, 1864, in Brady, Trans. Linn. Soc. London, Zool., v. 24, p. 471, pl. 48, fig. 18.
Elphidium arcticum (Parker and Jones), Cushman, 1930, U. S. Nat. Mus. Bull. 104, pt. 7, p. 27, pl. 11, figs. 1-6.
Elphidiella arctica (Parker and Jones), Cushman, 1939, U. S. Geol. Surv. Prof. Pap. 191, p. 65, pl. 18, figs. 11-14.
- Elphidium aculeatum* (d'Orbigny)
Polystomella aculeata d'Orbigny (sit.), 1846, Foraminiferes fossiles du Bassin Tertiaire de Vienne (Autriche). Gide et Comp. p. 131, pl. 6, figs. 27-28.
Elphidium aculeatum (d'Orbigny), Thalmann, 1932, Ecologiae Geologicae Helvetiae, v. 25, 293-312.
- Elphidium advena* (Cushman)
Polystomella advena Cushman, 1922, Carnegie Inst. Washington, Publ. 311, p. 56, pl. 9, figs. 11, 12.
Elphidium advenum (sit.), Cushman, 1930, U.S. Nat. Mus., Bull. 104, pt. 7, p. 25, pl. 10, figs. 1, 2.
- Elphidium articulatum* (d'Orbigny)
Polystomella articulata d'Orbigny, 1839, Amer. Merid., Foraminifères, v. 5, pt. 5, p. 30, pl. 3, figs. 9, 10.
Elphidium articulatum (d'Orbigny), Cushman, 1930, U.S. Nat. Mus., Bull. 104, pt. 7, p. 26, pl. 10, figs. 6-8.
- Elphidium bartletti* Cushman
Elphidium bartletti Cushman, 1933, Smithsonian Misc. Coll., v. 89, no. 9., p. 4, pl. 1, figs. 9a, b.

Foraminifera from the Kuromatsunai and Setana Formations

- Elphidium bartletti* Cushman, Loeblich and Tappan, 1953, Smithsonian Misc. Coll., v. 121, no. 7, p. 96, pl. 18, figs. 10-14.
- Elphidium crispum* (Linné)
Nautilus crispum Linné, 1758, Syst. Nat., ed. 10, p. 709.
Elphidium crispum (Linné), Cushman and Grant, 1927, San Diego Soc. Nat. Hist. Trans., v. 5, no. 6, p. 73, pl. 7, figs. 8a, b.
- Elphidium excavatum* (Terquem) forma *clavata* Cushman
Elphidium incertum (Williamson) var. *clavatum* Cushman, 1930, U. S. Nat. Mus., Bull. 104, pt. 7, p. 20, pl. 7, figs. 10a, b.
Elphidium decipiens (Costa), Hada, 1931, Tohoku Imp. Univ. Sci., Rep., 4th, ser. v. 6, no. 3, p. 126, text-figs. 83a, b.
Elphidium hughesi foraminosum Cushman, Asano, 1950, Illust. Cat. Japan. Tert. Small. Foram., pt. 1, p. 8, text-figs. 46, 47.
Elphidium clavatum Cushman, Loeblich and Tappan, 1953, Smith. Misc. Coll., v. 121, no. 7, p. 98, pl. 19, figs. 8-10.
Elphidium excavatum (Terquem) forma *clavata* Cushman, Takayanagi, 1955, Tohoku Univ., Inst., Geol., Pal., Contr., no. 45, p. 42, pl. 1, fig. 25.
- Elphidium frigidum* Cushman
Elphidium frigidum Cushman, 1933, Smithsonian Misc. Coll., vol. 89, no. 9, p. 5, pl. 1, fig. 8.
- Elphidium hanzawai* Asano
Elphidium hanzawai Asano, 1939, Jour. Geol., Soc., Japan. v. 46, no. 551, 426, figs. 3, 4a, b.
- Elphidium incertum* (Williamson)
Polystomella umbilicatula var. *incerta* Williamson, 1858, Recent Foraminifera of Great Britain, p. 44, pl. 3, fig. 82a.
Elphidium incertum (Williamson), Macfadyen, 1932, Geol. Mag., vol. 69, no. 821, pl. 35, figs. 6a, b.
- Elphidium jenseni* (Cushman)
Polystomella macella (Fichtel and Moll) var., Jensen, 1904, Linnean Soc. New South Wales Proc., v. 29, p. 817, pl. 23, fig. 4.
Polystomella jenseni Cushman, 1924, Carnegie Inst. Washington, Publ. 342, p. 49, pl. 16, figs. 4 (?), 6.
Elphidium jenseni (Cushman), Cushman, 1933, U.S. Nat. Mus., Bull. 161, pt. 2, p. 48, pl. 11, figs. 6, 7.
- Elphidium macellum* (Fichtel and Moll)
Elphidium macellus (Fichtel and Moll) var., 1798, Testacea microscopia, aliaque minuta ex generibus Argonauta et Nautilus, ad naturam picta et descripta. p. 68, pl. 10, figs. h, i, k.
Elphidium macellum var. *tumidocamerale* Bogdanowicz, 1932, Oil Geol., Inst., Leningrad, Trans., ser. A, fasc. 22, II, i, 2, text-figs. 9, 10.
Elphidium macellum (Fichtel and Moll), Cushman, 1939, Geol. Surv. prof. paper, 191, p. 51, pl. 14, figs. 1-3
- Elphidium subarcticum* Cushman
Elphidium subarcticum Cushman, 1944, Cushman Lab. Foram. Res., Spec., Publ., no.

- 12, p. 27. pl. 3, figs. 34, 35.
Elphidium nakanokawaense Shirai, 1960, Jour. Fac. Sci. Hokkaido Univ., Ser. IV, vol. X, 537-543, pl. 1, figs. 4a-b, 5a-b.
- Elphidium subgranulosum* Asano
Elphidium subgranulosum Asano, 1938, Jour. Geol. Soc. Japan, 45, p. 586, pl. 14, figs. 4a, b.
- Elphidium subincertum* Asano
Elphidium subincertum Asano, 1951, Illust. Cat. Japan Tert. Small Foram., pt. 1, p. 10, figs. 56, 57.
- Elphidium translucens* Natland
Elphidium translucens Natland, 1938, Scripps Inst. Oceanogr. Bull., Tech. ser., v. 4, p. 144, pl. 5, figs. 3, 4.
- Epistominella pulchella* Husezima and Maruhasi
Epistominella pulchella Husezima and Maruhasi, 1944, Jour. Shigenkagaku Kenkyusho (Res. Inst. Nat. Resour., Japan), v. 1, no. 3, p. 398, pl. 34, figs. 10a-c.
- Fissurina annectens* (Burrows and Holland)
Lagena annectens Burrows and Holland, 1895, in Jones, Parker and Brady, 1985, Palaeontogr. Soc., London, Part II, p. 203, pt. vii, II.
Fissurina annectens (Burrows and Holland), Barker, 1960, Spc. Publ., Soc. Economic Paleontologists and Mineralogists, 9.
- Fissurina baccata* (Hellon-Allen and Earland)
Lagena orbignyana var. *baccata* Hellon-Allen and Earland, 1922, Terra Nova Exped., Zool., VI, no. 2, p. 162, pt. vi, figs. 15, 16.
Fissurina baccata (Hellon-Allen and Earland), Jones, 1994, Challenger Foram., Pl. 59, Fig. 20.
- Fissurina lacunata* (Burrows and Holland)
Lagena castrensis Brady (not of Schwager), 1884, Voy. Challenger, Rep. Zool., v. 9, p. 485, pl. 60, figs. 1, 2.
Lagena lacunata Burrows and Holland, 1895, in Jones, T.R., Palaeontogr. Soc. London, p. 205, pl. 7, fig. 12.
Fissurina orbignyana lacunata (Burrows and Holland), Asano, 1938, Tohoku Imp. Univ., Sci. Rep., 2nd ser. (Geol.), v. 19, no. 2, p. 219, pl. 27, figs. 27, 28.
- Fissurina lucida* (Williamson)
Entosolenia marginata (Montagu) var. *lucida* Williamson, 1848, Ann. Mag. Nat. Hist., ser. 2, v. 1, p. 17, pl. 2, fig. 17.
Fissurina lucida (Williamson), Loeblich and Tappan, 1953, Smithsonian Misc. Coll., v. 121, no. 7, p. 76, pl. 14, fig. 4.
- Fissurina marginata* (Montagu)
Vermiculum marginatum Montagu, 1803, Testacea Britannica, p. 524.
Fissurina marginata (Montagu), Loeblich and Tappan, 1953, Smithsonian Misc. Coll., v. 121, no. 7, p. 77, pl. 14, figs. 6-9.
- Fissurina obscurocostata* Galloway and Wissler
Fissurina obscurocostata Galloway and Wissler, 1927, Jour. Pal., v. 1, p. 52, pl. 9, fig. 1.

Foraminifera from the Kuromatsunai and Setana Formations

Fissurina orbignyana Seguenza

Fissurina orbignyana Seguenza, 1862, *Foram. monotal. Mioc. Messina*, p. 66, pl. 2, figs. 24, 26.

Fissurina cf. *rizzae* Seguenza

cf. *Fissurina rizzae* Seguenza, 1862, *Foram. monotal.*, *Mioc. Messina*, p. 72, pl. 2, fig. 50.

Fissurina semimarginata (Reuss)

Lagena sp. (Nos. 64-65) von Schlicht, 1870. *Die Foraminiferen Septarienthones Pielzpuhl*, p. 11, pl. 4, figs. 4-6, 10-12.

Lagena marginata Williamson var. *semimarginata* Reuss, 1870, *Sitzb. Akad. Wiss. Wien*, v. 62, pt. 1, p. 468.

Fissurina semimarginata (Reuss), Barker, 1960, *Sp. Publ., Soc. Economic Paleontologists and Mineralogists*, 9.

Fissurina cf. *subquadrata* Parr

cf. *Fissurina subquadrata* Parr, 1954, *Roy. Soc. Victoria, Proc.*, n. s., LVI, pt. 2, 203, ix, 5.

Gavelinopsis praegeri (Heron-Allen and Earland)

Discorbina praegeri Heron-Allen and Earland, 1913, *Royal Irish Acad., Proc.*, v. 31, pt. 64, p. 122, pl. 10, figs. 8-10.

Gavelileopsis praegeri (Heron-Allen and Earland), Hofker, 1951, *Siboga Exped., Mon.* IV, pt. 3, p. 485, figs. 332-334.

Glabratella cf. *aurantista* Seiglie and Bermúdez

cf. *Glabratella aurantista* Seiglie and Bermúdez, 1965, *Monographia de la familia de foraminiferos Glabratellidae*, *Geos*, 1965, no. 12,

Glabratella mirabiris panamensis Seiglie and Bermúdez

Glabratella mirabiris panamensis Seiglie and Bermúdez, 1965, *Geos*, 12, p. 33, pl. 6, a-b, 7, 8.

Glabratella pulvinata (Brady)

Discorbina pulvinata Brady, 1884, *Rept. on the Sci. Reslt. of the Voy. of the H. M. S. Challenger during the years 1873-1876, Zoology*, v. 9,

Glabratella pulvinata, (Brady), Barker, 1960, *SEPM Spec. Publ.*, 9, pl. 88, figs. 10a, b.

Glabratella sp.A

This species is somewhat resembles *G. crassa* Hofker but differs from it in having more rounded and inflated chambers, and five to six chambers in final whorl.

Glabratella sp.B

This species is characterized in having planoconvex test with low trochospiral coiling and with long spines in its periphery.

Glandulina ovula d'Orbigny

Glandulina ovula d'Orbigny, 1846, *Foram. Foss. Bass. Tert. Vienne*, p. 29, pl. 1, figs. 6, 7.

Globocassidulina bisecta Nomura

Globocassidulina bisecta Nomura, 1983, *Tohoku Univ. Sci. Rep.*, 2nd ser. (Geol), v. 53, no. 1, p. 73-77, pl. 2, figs. 2,3; pl. 14, figs. 8-12; pl. 15, figs. 1-5.

Globocassidulina jamesoni (McCulloch)

Cassidulina (?) *jamesoni* McCulloch, 1977, *Quantitative observations on Recent foram. tests, with emphasis on the eastern Pacific*, p. 390, pl. 164, figs. 15a-c.

- Globocassidulina jamesoni* (McCulloch), Nomura, 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 54, no. 1, p. 33, 34, pl. 3, figs. 7a-c, 8a, b; pl. 19, figs. 1, 2.
- Globocassidulina canalisuturata* Eade
Globocassidulina canalisuturata Eade, 1967, N.Z. Jour. Mar. Freshwater Res., 1. no. 4, p. 440, fig. 3 (5-7), fig. 5 (7, 8).
- Globocassidulina neobrecha* Nomura
Globocassidulina neobrecha Nomura, 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 53, no. 1, p. 67-68, pl. 1, figs. 16a-c; pl. 19, figs. 3-6.
- Guttulina yabei* Cushman and Ozawa
Guttulina yabei Cushman and Ozawa, 1929, Japan Jour. Geol. Geogr., v. 6, nos. 3-4, p. 68, pl. 13, fig. 2, pl. 14, fig. 6.
- Hanzawaia nipponica* Asano
Hanzawaia nipponica Asano, 1943, Jour. Geol. Soc. Japan, v. 51, no. 606, p. 98, pl. 4, figs. 1-2.
- Heterolepa praecincta* (Karrer)
Rotalia praecinctus Karrer, 1868, Sitz. Akad. Wiss. Wien, v. 58, p. 189, pl. 5, fig. 7.
Truncatulina praecincta Cushman, 1915, U.S. Nat. Mus. Bull., v. 71, no. 5, p. 39, pl. 26, fig. 2. *Eponides praecinctus* (Karrer), Asano, 1951, Illust. Cat. Japan Tert. Small Foram., pt. 14, p. 11, figs. 30-32.
- Heterolepa subhaidingerii* (Parr)
Cibicides subhaidingerii Parr, 1950, Foraminifera, Rep. B.A.N.Z. Antarctic Res. Exp. 1929-1931, Ser. B. (Zoology and Botany), 5, p. 364, pl. 15, fig. 7.
Heterolepa subhaidingerii (Parr), Tappan and Loeblich, 1982, Synopsis and Classification Living Organisms, pl. 53, fig. 10.
- Islandiella helenae* Feyling-Hanssen and Buzas
Cassidulina teretis Tappan, Loeblich and Tappan (not of Tappan, 1951), 1953, Smithsonian Misc. Coll., v. 121, no. 7, p. 121, pl. 24, figs. 3, 4.
Islandiella helenae Feyling-Hanssen and Buzas, 1976, Jour. Foram. Res., v. 6, no. 2, p. 155, figs. 1-4.
- Islandiella islandica* (Nørbang)
Cassidulina islandica Nørbang, 1945, Foraminifera. in the Zoology of Iceland. v.2, p. 42, text-figs. 7, 8d-f.
Islandiella islandica (Nørbang), Nørbang, 1958, Vidensk. Medd. Dansk Natur-hist. Foren., v. 120, p. 27, 28, pl. 6, figs. 1a, b, 2-5, pl. 7, figs. 6a, b, 7a, b.
- Islandiella japonica* (Asano and Nakamura)
Cassidulina japonica Asano and Nakamura, 1937, Japan Jour. Geol. Geogr., v. 14, nos. 3-4, p. 144, pl. 13, figs. 1, 2.
Islandiella japonica (Asano and Nakamura), Troitskaja, 1970, Usloviya obitaniya i raspredelenie foraminifer v Yaponskom More (semeistva Elphidiidae, Cassidulinidae i Islandiellidae). In Fursenko A.V. ed., Obstchie voprosy izucheniya mikrofauny Sibiri, dalinego vostoka i drugikh raionov. Instituta Geologii i Geofiziki, Akademiya Nauk SSSR, Sibirskoe Otdelenie, Trudy, vol. 71, p. 150, pl. 6, figs. 3, 4.
- Islandidlla norcrossi* (Cushman)

Foraminifera from the Kuromatsunai and Setana Formations

- Cassidulina norcrossi* Cushman, 1933, Smithsonian Misc. Coll., v. 89, no. 9, p. 7, pl. 2, figs. 7a-c.
- Islandiella norcrossi* (Cushman), Nørbang, 1958, Dansk Naturh. Foren. København Vidensk. Meddel., v. 120, p.26.
- Islandiella setanaensis* (Asano and Nakamura)
- Cassidulina setanaensis* Asano and Nakamura, 1937, Japan Jour. Geol. Geogr., v. 14, nos. 3-4, pl. 13, figs. 7a,b
- Islandiella setanaensis* (Asano and Nakamura), Nomura, 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 53, no. 1, p. 1-101, pl. 3, fig. 6, pl.4, figs. 3,4, pl. 10, fig. 11, pl.11, fig. 1-3; 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 54, no. 1, p. 5, pl. 1, figs. 3a-c.
- Islandiella sublimbata* (Asano and Nakamura)
- Cassidulina sublimbata* Asano and Nakamura, 1937, Japan Jour. Geol. Geogr., v. 14, nos. 3-4, p. 146, pl. 14, figs. 3, 4a, b.
- Islandiella sublimbata* (Asano and Nakamura), Nomura, 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 53, no. 1, pl. 3, fig. 7, pl. 4, fig. 2, pl. 5, fig. 7, pl. 9, figs. 1-8; 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 54, no. 1, p. 9, pl. 1, figs. 6-8.
- Islandiella yabei* (Asano and Nakamura)
- Cassidulina yabei* Asano and Nakamura. 1937, Japan. Jour. Geol. Geogr., v. 14, nos. 3-4, p. 145, pl. 14, figs. 1a, b.
- Islandiella yabei* (Asano and Nakamura), Nomura, 1983, Tohoku Univ. Sci. Rep., 2nd ser. (Geol), v. 53, no. 1, pl. 3, fig. 1; pl. 4, fig. 7; pl. 9, figs. 9-12, pl. 10, figs. 1-3.
- Karreriella baccata* (Schwager) *japonica* Asano
- Karreriella baccata* (Schwager) *japonica* Asano, 1938, Japanese Jour. Geol. Geogr., v.15, nos. 1-2, p. 90, pl. 10, figs. 1a-c.
- Lagena acuticosta* Reuss
- Lagena acuticosta* Reuss, 1861, Sitz. Akad. Wiss. Wier., v. 44, no. 1, p. 305, pl. 1, fig. 4.
- Lagena apiopleura* Loeblich and Tappan
- Lagena apiopleura* Loeblich and Tappan, 1953, Smithsonian Misc. Coll., v. 121, no. 7, p. 59, pl. 10, figs. 14, 15.
- Lagena striata* (d'Orbigny)
- Oolina striata* d'Orbigny, 1839, Voy. Amer. Merid., Foraminifères, v. 5, pt. 5, p. 21, pl. 5, fig. 12.
- Lagena striata* (d'Orbigny), Asano, 1938, Tohoku Imp. Univ., Sci. Rep., 2nd ser. (Geol.), v. 19, p. 217, pl. 27, fig. 26; pl. 28, fig. 28.
- Lagena sulcata spicata* Cushman and McCulloch
- Lagena sulcata* var. *spicata* Cushman and McCulloch, 1950, Allan Hancock Pacific Exped., v. 6, no. 6, p. 360, pl. 48, figs. 3-7.
- Lenticulina nikobarensis* (Schwager)
- Cristellaria nikobarensis* Schwager, 1886, Novara Exp. Geol. Thell., 2, p. 243, pl. 6, fig. 87.
- Robulus nikobarensis* (Schwager), Asano, 1938, Sci. Rep. Tohoku Univ. ser. 2, 19 (2), p. 204, pl. 28, figs. 5, 6; pl. 29, fig. 8.
- Lenticulina nikobarensis* (Schwager), Loeblich and Tappan, 1988, Foraminiferal genera and their classification. p. 405. Pl. 446, figs. 9, 10.

Melonis pompilioides (Fichtel and Moll)

Nautilus pompilioides Fichtel and Moll, 1798, Testacea microscopica, p. 31, pl. 2, figs. a-c.

Melonis etruscus Montfort, 1808, Conch. Syst., p. 67.

Melonis pompilioides (Fichtel and Moll), Voloshinova, 1958, Mikrofauna SSSR, Sbornik 9, VNIGRI, Trudy, p. 149, pl. 3, fig. 1

Melonis uchioi Hasegawa

Melonis uchioi Hasegawa, 1991, Trans. Proc. Palaeont. Soc. Japan, N.S., no. 164, p. 1003.

Melonis pacificus (Cushman)

Nonion umbilicatula var. *pacifica* Cushman, 1924, Carnegie Inst. Wash., v. 342, p. 48, pl. 16, fig. 3.

Nonion pacificum Cushman, 1939, U.S. Geol. Surv., Prof. paper v. 191, p.25, pl. 6, fig. 25.

Melonis pacificus (Cushman), Hasegawa, 1979, Tohoku Univ., Sci. Rep., 2nd ser. (Geol), v. 49, no. 2, p. 151.

Miliolinella circularis (Bornemann)

Triloculina circularis Bornemann, 1855, Zeit. deutsch. Geol. Ces., v. 7, p. 349, pl. 19, fig. 4

Miliolinella circularis (Bornemann), Asano, 1951, Illust. Cat. Japan. Tert. Small. Foram., pt. 6, p. 9, figs. 65-67.

Neoconorbina stachi (Asano)

Discopulvinulina stachi Asano, 1951, Illust. Cat. Japan Tert. Small. Foram., pt. 14, p. 7, figs. 46-48.

Neoconorbina stachi (Asano), Matoba, 1970, Tohoku Univ., Sci. Rep., 2nd ser. (Geol), v. 42, no. 2, p. 57, pl. 4, flgs. 6a-c.

Neoeponides procerus (Brady)

Pulvinulina procera Brady, 1884, Voy. Challenger, Rep., Zool., v. 9, p. 698, pl. 105, figs.

7a-c. *Neoeponides procerus* (Brady), Reiss, 1960. Israel Geol. Surv., Bull. no. 29, p. 17.

Nonion japonicus Asano

Nonion japonicum Asano (sic.) , 1938, Jour. Geol. Soc. Japan, v. 45, no. 593, pl. 15(4), figs. 1, 2.

Nonionella stella Cushman and Moyer

Nonionella miocenica Cushman var. *stella* Cushman and Moyer, 1930, Cushman Lab. Foram. Res., Contr., v. 6, pt. 1, p. 56, pl. 7, figs. 17a-c.

Nonionellina labradorica (Dawson)

Nonionina labradorica Dawson, 1860, Canad. Nat., v. 5, p. 191, text-fig. 4.

Nonionellina labradorica (Dawson), Voloshinova, 1958, Mikrofauna SSSR, Sbornik 9, VNIGRI, Trudy, no. 115, p. 142.

Nonionoides grateloupi (d'Orbigny)

Nonionina grateloupi d'Orbigny, 1826, Annales des Sciences Naturelles v. 7, p. 294,

Nonionoides grateloupi (d'Orbigny), Saidova, 1975, BENTOSNYE FORAMINIFERY TIKHOGO Okeana, 3 vol., p. 248.

Oolina carteri Albani and Yassini

Oolina carteri Albani and Yassini, 1989, Aust. J. Mar. Freshw. Res., v. 70, p. 385, fig. 3r-s.

Oolina costata (Williamson)

Entosolenia costata Williamson, 1858, Recent Foraminifera Great Britain, p. 9, pl. 1, fig.

18.

Lagena costata (Williamson), Cushman, 1923, U.S. Nat. Mus. Bull. v. 104, pt.4, p.12, pl.1, fig. 16, pl. 2, figs. 1, 2 (not pl. 3, fig. 8)

Oolina costata (Williamson), Parker, 1952, Bull. Mus. Comp. Zool., v. 106, no. 9, p. 409, pl. 4, figs. 20,21.

Oolina melo d'Orbigny

Oolina melo d'Orbigny, 1839, Voy. Amer. Merid., Foraminifères, v. 5, pt. 5, p. 20, pl. 5, fig. 9.

Oolina striatopunctata (Parker and Jones)

Lagena sulcata (Walker and Jacob) var. *striatopunctata* Parker and Jones, 1865, Philos. Trans. Roy. Soc. London, v. 155, p. 350, pl. 13, figs. 25-27.

Entosolenia striatopunctata (Parker and Jones) Dawson, 1870, Can. Nat., n.s., v. 5, p. 178, fig. 11. *Lagena striatopunctata* Parker and Jones, Brady, 1878, Ann. Mag. Nat. Hist., ser. 5, v. 1, p. 434, pl. 20, fig. 3.

Oolina striatopunctata (Parker and Jones) Loeblich and Tappan, 1953, Smithsonian Misc. Coll. v. 121, no. 7, p. 74, pl. 12, figs. 2-5.

Oridorsalis umbonatus (Reuss)

Rotalia umbonata Reuss, 1851, Deutsch. geol. Gesell., Zeitschr., v. 3, p. 75, pl. 5, figs. 35a-c.

Oridorsalis umbonatus (Reuss), Parker, 1964, Jour. Pal., v. 38, no. 4, p. 626, pl. 99, figs. 4-6.

Osangulariella umbonifera (Cushman)

Pulvinulinella umbonifera Cushman, 1933, Cushman Lab. Foram. Res., Contr., 9, 90.

Osangulariella umbonifera (Cushman), Jones, 1994, Challenger Foraminifera, p. 99, pl. 95, figs. 9, 10.

Paracassidulina sulcata Belford

Paracassidulina sulcata Belford, 1966, Bur. Min. Resour. Aust. Rep., no. 79, p. 142, 144, pl. 24, figs. 11-14, text-fig. 16, nos. 7, 8.

Pararotalia nipponica (Asano)

Rotalia nipponica Asano, 1936, Jour. Soc. Japan, v. 43, no. 515, p. 614, pl. 30, figs. 2a-c.

Pararotalia nipponica (Asano), Ujiie, 1966, Palaeont. Soc. Japan, Trans. Proc., N.S.m no. 61, p. 192, text-figs. 1-3; pl. 24, figs. 1-7; pl. 25, figs. 1-5.

Patellina corrugata Williamson

Patellina corrugata Williamson, 1858, Rec. Foram. Gt. Britain, p. 46, pl. 3, figs. 86-89.

Patellinella hanzawai Asano

Patellinella hanzawai Asano, 1936, Jour. Geol. Soc. Japan, v. 43, no. 515, p. 613, pl. 31, figs. 3a-c.

Pateoris hauerinoides (Rhumbler)

Quinqueloculina subrotunda (Montagu) forma *hauerinoides* Rhumbler, 1936, Foram. der Kieler Bucht, Teil II-Ammodisculinidae bis Textulinidae, v. 1, no. 1, p. 206, 217, 226, text-figs. 167, 208-212.

Pateoris hauerinoides (Rhumbler) Loeblich and Tappan, 1953, Smithsonian, Misc. Coll. v. 121, no. 7, p. 42, pl. 6, figs. 8-12, text-figs. 1A, B.

Planoglabratella australensis (Heron-Allen and Earland)

Discorbis australensis Heron-Allen and Earland, 1932, *Discovery Repts.*, IV, 416.

Glabratella australensis (Heron-Allen and Earland), Loeblich and Tappan, 1964, *Treatise on Invertebrate*. p. 588.

Planoglabratella opercularis (d'Orbigny)

Rosalina opercularis d'Orbigny, 1839, *Foraminifères*, p. 93, pl. 3, figs. 24, 25.

Discorbis opercularis Cushman, 1915, *U.S. Nat. Mus. Bull.*, vol. 71, no. 5, p. 18, pl. 11, fig. 3.

Planolabratella opercularis (d'Orbigny), Seiglie and Bermúdez, 1965, *Geos*, 1965, no. 12, 15-65.

Planoglabratella patelliformis (Brady)

Discorbina patelliformis Brady, 1884, *Rept. on the Sci. Reslt. of the Voy. of the H. M. S. Challenger during the years 1873-1876, Zoology*, v. 9,

Glabratella patelliformis (Brady), Seiglie and Bermúdez, 1965, *Monographia de la familia de foraminíferos Glabratellidae*, *Geos*, 1965, no. 12,

Planoglabratella subopercularis (Asano)

Discorbis subopercularis Asano, 1951, *Illust. Cat. Japan Tert. Small Foram.*, pt. 14, p. 3, figs. 17-19.

Glabratella subopercularis (Asano), Matoba, 1970, *Tohoku Univ., Sci. Rep.*, 2nd ser. (Geol.), v. 42, no. 1, p. 54, pl. 5, figs. Sa-c.

Planolabratella subopercularis (d'Orbigny), Seiglie and Bermúdez, 1965, *Geos*, 1965, no. 12, 15-65.

Planoglabratella ? sp. C

This species resembles *Planoglabratella subopercularis* (Asano). but differs from it in havng less number of chambers in final whorl.

Polystomelina discorbinoides Yabe and Hanzawa

Polystomelina discorbinoides Yabe and Hanzawa, 1923, *Japan Jour. Geol. Geogr.*, v. 2, p. 99, text-flgs. a-c.

Poroeponides cribrorebandus Asano and Uchio

Poroeponides cribrorebandus Asano and Uchio, 1951, *Illust. Cat. Japan. Tert. Small. Foram.*, pt. 14, p.18, figs. 134, 135.

Pseudononion japonicum Asano

Pseudononion japonicum Asano, 1936, *Jour. Geol. Soc. Japan*, v. 43, no. 512, p. 347, text-figs. a-c.

Pseudoparrella naraensis Kuwano

Pseudoparrella naraensis Kuwano, 1950, *Jour. Geol. Soc. Japan*, v. 56, no. 657, p. 317, text-figs. 6a-c.

Pseudoparrella takayanagii (Iwasa)

Epistominella takayanagii Iwasa, 1955, *Jour. Geol. Soc. Japan*, v. 61, no. 712, p. 16, text-figs. 4a-c.

Epistominella suttsuensis Shirai, 1960, *Jour. Fac. Sci. Hokkaido Univ.*, Ser. IV, vol X, 537-543, pl. 2, figs. 3a-c.

Pseudoparrella takayanagii (Iwasa), Hasegawa, 1979, *Tohoku Univ., Sci. Rep.*, 2nd ser.

Foraminifera from the Kuromatsunai and Setana Formations

- (Geol), v. 49, no. 2, p. 153, pl. 5, figs. 7a-c.
- Pseudopolymorphina suboblonga* Cushman and Ozawa
Pseudopolymorphina suboblonga Cushman and Ozawa, 1930, U.S. Nat. Mus., Proc., v. 77, p. 91, pl. 23, figs. 3a-c.
- Pullenia apertura* Cushman
Pullenia apertura Asano (not of Cushman, 1927), 1951, Illust. Cat. Japan. Tert. Small. Foram., pt. 12, p. 10, figs. 3, 4.
- Pyrgo ezo* Asano
Pyrgo ezo Asano, 1938, Japanese Jour. Geol. Geogr., v.15, nos. 1-2, p.93, pl. 9, figs. 1-4, 6.
- Pyrgo fomasinii* Chapman and Parr
Pyrgo fomasinii Chapman and Parr, 1935, Roy. Soc. W. Australia, Jour., XXI, p. 5.
- Quinqueloculiua agglutinata* Cushman
Quinqueloculiua agglutinata Cushman, 1917, U.S. Nat. Mus., Bull. 71, pt. 6, p. 43, pl. 9, figs. 2a-c.
- Quinqueloculina akneriana* d'Orbigny
Quinqueloculina akneriana d'Orbigny, 1846, Foram. Foss. Bas. Vienne, p. 290, pl. 18, figs. 16-21.
- Quinqueloculina costata* d'Orbigny
Quinqueloculina costata d'Orbigny, 1826, Ann. Sci. Nat., v. 7, p. 301. no. 3.
- Quinqueloculina elongata* Natland
Quinqueloculina elongata Natland, 1938, Bull. Scripps Inst. Oceanogr. Tech. Ser., v. 4, no.5, p. 141, pl. 4, fig. 5.
- Quinqueloculina kuromatunaiensis* Asano
Quinqueloculina kuromatunaiensis Asano, 1936, Jour. Geol. Soc. Japan, v. 43, no. 515, p. 621, pl. 32, figs. 4a-d.
- Quinqueloculina sawanensis* Asano
Quinqueloculina sawanensis Asano, 1951, Illust. Cat. Japan. Tert. Small. Foram., pt. 6, p. 6, figs. 40-42.
- Quinqueloculina seminulum* (Linné)
Serpula seminulum Linné, 1758, Syst. Nat., ed. 10, p. 786.
Quinqueloculina seminulum (Linné), d'Orbigny, 1826, Ann. Sci. Nat., Paris, ser. 1, v. 7, . 303, no. 4.
- Quinqueloculina vulgaris* d'Orbigny
Quinqueloculina vulgaris d'Orbigny, 1826, Ann. Sci. Nat., v. 7, p. 302, no. 33.
- Quinqueloculina* cf. *vulgaris* d'Orbigny
cf. *Quinqueloculina vulgaris* d'Orbigny , 1826, Ann. Sci. Nat., v. 7, p. 302, no. 33.
- Quinqueloculina yessoensis* Asano
Quinqueloculina yessoensis Asano, 1936, Jour. Geol. Soc. Japan. 43, p. 620, pl. 32, figs. 3a-c.
- Quinqueloculina* sp.B

This species is somewhat similar to *Quinqueloculina elongata* Natland of Asano (1950) in its outline, but is distinguished from it by more inflated chambers and coarsely perforated wall.

Quinqueloculina sp.C

This species is somewhat similar to *Quinqueloculina contrita* d'Orbigny, but is distinguished by more rounded outline, short size, and apertural view.

Quinqueloculina sp.F

This form is similar to *Quinqueloculina yessoensis* Asano, but it differs from it in having a broadly oval test with more sharply angled periphery in apertural view. This is also similar to *Quinqueloculina kuromatunaiensis* Asano, but is distinguished in having angled periphery of chambers.

Quinqueloculina sp.G

This form is somewhat similar to *Quinqueloculina elongata* Natland, but is distinguished from it by its triangular appearance in apertural view.

Quinqueloculina sp.H

This form is similar to *Quinqueloculina hasimotoi* Asano, but differs from it in having a test without apertural protrusion.

Rectobolivina raphanuss (Parker and Jones)

Uvigerina (*Sagrina*) *raphana* Parker and Jones, 1865, Roy. Soc. London Philos. Trans, v. 155, p. 364, pl. 18, figs. 16, 17.

Rectobolivina raphana (sic.) (Parker and Jones), Loeblich and Tappan, 1964, Treatise on Invertebrate Pal., Moore, R.C., ed., pt. C, Protista 2, v. 2, p. C533, figs. 438 (9-11).

Rosalina australis (Parr)

Discorbis australis Parr, 1932, Roy. Soc. Victoria, Proc., XLIV (n.s.) p. 227, pl. xxii, 31.

Rosalina australis (Parr), Jones, 1994, Challenger Foram., pl. 87, figs. 5-7.

Rosalina bradyi (Cushman)

Discorbina globularis Brady (not of d'Orbigny, 1826), 1884, Voy. Challenger, Rep., Zool., v. 9, p. 86, figs. 8a-c.

Discorbina globularis var. *bradyi* Cushman, 1915, U.S. Nat. Mus., Bull. 71, pt. 5, p. 12, pl. 8, figs. 1a-c.

Rosalina bradyi (Cushman), Hornibrook and Vella, 1954, Micropal., v. 8, no. 1, p. 26.

Rosalina globularis d'Orbigny

Rosalina globularis d'Orbigny, 1926, Ann. Sci. Nat., Paris, ser. 1, v. 7, p. 271, no. 1, pl. 13, figs. 1-14; Modele no. 69.

Rosalina isabelleana d'Orbigny

Rosalina isabelleana d'Orbigny, 1839, Foraminifères. p. 43, pl. 6, figs. 10-12.

Rosalina vilardeboana d'Orbigny

Rosalina vilardeboana d'Orbigny, 1839, Voy. Amer. Merid., Foraminifères, v. 5, pt. 5, p. 44, pl. 6, figs. 13-15.

Rosalina vilardeboana d'Orbigny

Rosalina vilardeboana d'Orbigny, 1839, Voy. Amér. Mérid., Foraminifères, v. 5, pt. 5, p. 44, pl. 6, figs. 13-15.

Sigmoidella pacifica Cushman and Ozawa

Sigmoidella (*Sigmoidina*) *pacifica* Cushman and Ozawa, 1928, Cushman Lab. Foram. Res., Contr., v. 4, p. 19, pl. 2, fig. 13.

Sigmoilina sigmoidea (Brady) *compressa* Cushman

Foraminifera from the Kuromatsunai and Setana Formations

- Sigmoilina sigmoidea* (Brady) *compressa* Cushman, 1946, Contr. Cushman Lab. Foraminifera Res., 22, no. 2, p. 32, pl. 5, figs. 10-12.
- Sigmomorphina semitecta terquemiana* (Fornasini)
Polymorphina amygdaloides Reuss var. *terquemiana* Fornasini, 1902, Mem. Accad. Inst. Sci. Bologna, ser. 5, no. 9, p. 72, fig. 25.
Sigmomorphina semitecta (Reuss) var. *terquemiana* Cushman and Ozawa, 1930, U.S. Nat. Mus. Proc., v. 77, no. 6, p. 129, pl. 33, figs. 4, 5; pl. 34, figs. 2, 3; pl. 35, fig. 1.
- Sigmomorphina trilocularis* (Bagg)
Polymorphina trilocularis Bagg, 1912, U. S. Geol. Surv. Bull., 513, p. 75, pl. 20, figs. 15-18.
Sigmomorphina trilocularis (Bagg), Cushman and Ozawa, 1930, U. S. Nat. Mus. Proc., v. 77, no. 6, p. 136, pl. 36, fig. 5.
- Sphaeroidina japonica* Asano
Sphaeroidina japonica Asano, 1953, Tohoku Univ., Inst. Geol. Paleont., Short Papers, no. 5, p. 17, pl. 2, figs. 43, 44.
- Spirillina limbata* Brady
Spirillina limbata Brady, 1879, Quart. Jour. Microscopic Sci., n.s., vol. 19, p. 64, pl. 8, figs. 26a, b.
- Spirillina vivipara* Ehrenberg
Spirillina vivipara Ehrenberg, 1843, Phys. Ab. K. Akad. Wiss. Berlin, 1841 (1), p. 323, 422, pl. 3, VII, fig. 41.
- Spiroloculina hadai* Thalmann
Spiroloculina costata Hada, 1931, Tohoku Univ. Sci. Rep., 4th ser., Biol., v. 6, no. 1, p. 84, text-figs. 37a,b.
Spiroloculina hadai Thalmann, 1933, Jour. Paleont., v. 7, no. 30, p. 354.
- Spiroplectammina* sp.A
This form somewhat similar to *Textularia candèiana* d'Orbigny, but has a test with planispirally arranged chambers in the early portion.
- Spiroplectinella wrightii* (Silvestri)
Spiroplecta wrightii Silvestri, 1903, Accad. Pont. Nuovi Lincei, Atti, LVI, p. 59, pl. 1x, figs. 1-6.
- Textularia conica* d'Orbigny
Textularia conica d'Orbigny, 1839, Foraminifères, p. 143, pl. 1, figs. 19, 18.
- Triloculina rotunda* d'Orbigny
Triloculina rotunda d'Orbigny, 1826, Ann. Sci. Nat. v. 7, p. 299, no. 4.
- Triloculina suttuensis* Asano
Triloculina suttuensis Asano, 1936, Jour. Geol. Soc. Japan, v. 43, no. 515, p. 621, pl. 33, figs. 2a-c.
- Triloculina tricarinata* d'Orbigny
Triloculina tricarinata d'Orbigny, 1826, Ann. Sci. Nat., 7, p. 299, no. 7.
- Uvigerina akitaensis* Asano
Uvigerina akitaensis Asano, 1950, Illust. Cat. Japan. Tert. Small. Foraminifera, pt. 2, p. 14, figs. 60-62.

Uvigerina juncea Cushman and Todd

Uvigerina juncea Cushman and Todd, 1941, Cushman Lab. Foram. Res., Contr., XVII, p. 78, pl. xx, 4-11.

Uvigerina schencki Asano

Uvigerina schencki Asano, 1950, Illust. Cat. Japan. Tert. Small. Foram. pt. 2, p. 17, figs. 74, 75.

Valvulineria hamanakoensis (Ishiwada)

Anomalina hamanakoensis Ishiwada, 1958, Geol. Surv. Japan Rep., no. 180, p. 18, text-figs. 3a-c; pl. 1, 24-27.

Valvulineria hamanakoensis (Ishiwada), Matoba, 1970, Tohoku Univ., Sci. Rep., 2nd ser. (Geol.), v. 42, no. 1, p. 63, pl. 4, figs. 12, 13.

Valvulineria japonica Asano

Valvulineria japonica Asano, 1951, Illust. Cat. Japan. Tert. Small. Foram. pt. 14, p. 7, figs. 49-51.

Acknowledgments

The author would like to express his deep gratitude to Associate Professor Shiro Hasegawa of the Laboratory of Geosphere Science, Graduate School of Environmental Earth Science, Hokkaido University, for his encouragement and advice during the course of this study. The author also wishes to thank Professor Yoshihiro Togo and Associate Professor Akihiko Suzuki of the Department of Earth Science, Iwamizawa College, Hokkaido University of Education, for their suggestions concerning various aspects of the present study. Sincere thanks are due to Dr. Yoshinori Hikida of Nakagawa Museum, Mr. Hiroyuki Takata, Mr. Takuya Itaki, and Mr. Jun'ichi Uchida of the Laboratory of Geosphere Science, Graduate School of Environmental Earth Science, Hokkaido University for their encouragement, and help in collecting references.

Deep appreciation is expressed to Mr. Yasuhumi Terasaki, the chief curator of the Imakane Board of Education, for his kind intentions during the course of this study.

References

- Asano K., 1936a, Foraminifera from Kuromatsunai-mura, Suttu-gun, Hokkaido. Studies on the fossil Foraminifera from the Neogene of Japan. (II). Jour. Geol. Soc. Japan, 43, 419-422. (in Japanese)
- Asano K., 1936b, Foraminifera from Kuromatsunai-mura, Suttu-gun, Hokkaido. Studies on the fossil Foraminifera from the Neogene of Japan. Part 2. Jour. Geol. Soc. Japan, 43, 615-624.
- Asano K., and Nakamura M., 1937, On the distribution of the Japanese species of Cassidulina. Jour. Geol. Soc. Japan, 44, 42-49. (in Japanese)
- Asano K., 1937, Pliocene foraminifera from the Setana Beds, Hokkaido. Jour. Geol. Soc. Japan, 44, 509-512. (in Japanese)

Foraminifera from the Kuromatsunai and Setana Formations

- Asano K., 1939, Pliocene (Upper Mizuho) Foraminifera from Japan (preliminary Report). Jour. Geol. Soc. Japan, 46, 155-168. (in Japanese with English abstract)
- Hashimoto W., Kanno S., Shinada Y., and Oshima K. , 1963, Geology of the Imagane, Kun'nui, and Yakumo districts, Oshima Peninsula, Hokkaido. Jour. Geol. Soc. Japan, 69, 228-238. (in Japanese with English abstract)
- Kato M., Katsui Y., Kitagawa Y., and Matsui M. (eds), 1990, Regional geology of Japan. Part 1. Hokkaido. Kyoritsu Shuppan, 337p.
- Masatani K., and Ohkura T., 1980MS, Neogene biostratigraphy in Oshima Peninsula, Hokkaido, Japan. In particular relation between "*Operculina-Miogypsina* Zone" and Planktonic foraminifera. Technol. Lab. JAPEX, 23, 32-52. (in Japanese with English abstract)
- Matsui M., Doi S., Takeda H., Uozumi S., Hujie T., Akiba C., Yoshimura N., Yamaguchi K., Obara T., and Odagiri T. , 1955, Report on geology and ore depositions in the north-eastern Imakane area, Shiribeshi, Hokkaido, Japan. Imakane town, 56. (in Japanese)
- Nagao T., and Sasa Y., 1933, The Cenozoic systems of southwestern Hokkaido and its geological history, part 1-2. Jour. Geol. Soc. Japan, 40 555-577; 750-775. (in Japanese)
- Nojo A., Togo, Y., Suzuki, A. and Sagayama, T., 1994, Lithostratigraphy and chronology of the Neogene Kuromatsunai Formation in the Imakane area, southwestern Hokkaido, Japan. Jour. Geol. Soc. Japan, 100, 771-786. (in Japanese with English abstract)
- Nojo A., Uchida J., Soeda Y., Itagaki K., Nakagawa M., Nikaido A., Suzuki A. and Togo Y., 1997, Some Plio-Pleistocene foraminifera from the Imakane-Kamiyakumo area of southwestern Hokkaido, Japan (Preliminary report). Bulletin of the studies in Imakane district, no. 3, 29-37. (in Japanese)
- Nojo A., Hasegawa S., Okada H., Togo Y., Suzuki A., and Matsuda T., in press, Interregional lithostratigraphy and biostratigraphy of the Pleistocene Setana Formation, southwestern Hokkaido, Japan. Geol. Soc. Japan. (in Japanese with English abstract)
- Oka T., and Mitani K. , 1981, Geology of the Imakane town. Imakane town, 77. (in Japanese)
- Shirai T., 1959, Geology in the Kuromatsunai district, with a special reference to the fossil foraminifera of the Pliocene Setana Formation—A study of the Neogene fossil foraminifera in northern Japan (1). Bull. Geol. Comm. Hokkaido, 38, 1-12. (in Japanese with English abstract)
- Shirai T., 1960, New genus and species of foraminifera from the Pliocene formation, southwestern Hokkaido. Jour. Fac. Sci. Hokkaido Univ., Ser. 4, 10, 537-543.

Plates (Scale bar = 100 μ m)

Planktonic Foraminifera (Plate 1)

1. *Globigerina bulloides* d'Orbigny, 2. *Globigerina quinqueloba* Natland,
3. *Globigerinita uvula* (Ehrenberg), 4. *Globigerinita glutinata* (Egger),
5. *Globorotalia inflata* (d'Orbigny), 6. *Globigerinoides ruber* (d'Orbigny),
- 7-9. *Neogloboquadrina incompta* (Chifelli), 10-11. *Neogloboquadrina pachyderma* (Ehrenberg)

Benthic Foraminifera (Plate 2-12)

Plate 2

1. *Angulogerina kokozuraensis* (Asano), 2. *Uvigerina akitaenesis* Asano,
3. *Uvigerina schencki* Asano, 4. *Uvigerina juncea* Cushman and Todd,
5. *Rectobolivina rephanus* (Parker and Jones), 6. *Buliminella elegantissima* (d'Orbigny),
7. *Bolivinita quadrilatera* (Schwager), 8. *Spiroplectammina* sp. A, 9. *Bolivina* sp. A,
10. *Bolivina decussata* Brady, 11. *Spirillina limbata* Brady,
12. *Cancris auriculus* (Fichtel and Moll)

Plate 3

1. *Oolina melo* d'Orbigny, 2-3. *Lagena apiopleura* Loeblich and Tappan,
4. *Oolina costata* (Williamson), 5. *Fissurina lacunata* (Burrows and Holland),
6. *Lagena striata* (d'Orbigny), 7. *Oolina striatopunctata* (Parker and Jones),
8. *Lagena acuticosta* Reuss, 9. *Fissurina baccata* (Hellon-Allen and Earland),
10. *Fissurina orbignyana* Seguenza, 11. *Fissurina* cf. *rizae* Seguenza,
12. *Fissurina annectens* (Burrows and Holland), 13. *Fissurina* cf. *subquadrata* Parr,
14. *Fissurina marginata* (Montagu)

Plate 4

1. *Cibicides lobatulus* (Warker and Jacob), 2-3. *Cibicides refulgens* de Montfort,
- 4-5. *Cibicides subdepressus* (Asano), 6. *Heterolepa subhaidingerii* (Parr),
7. *Anomalinoides globulosus* (Cushman and Parr),
8. *Dyocibicides perforata* Cushman and Valentine

Plate 5

1. *Buccella frigida* (Cushman), 2. *Buccella makiyamai* (Chiji), 3. *Buccella tenerrima* (Brady),
4. *Patellinella hanzawai* Asano, 5. *Hanzawaia nipponica* Asano, 6. *Rosalina australis* (Parr),

Foraminifera from the Kuromatsunai and Setana Formations

7. *Rosalina bradyi* (Cushman), 8. *Guttulina yabei* Cushman and Ozawa

Plate 6

1. *Elphidium subgranulosum* Asano, 2. *Elphidium frigidum* Cushman,
3-5. *Elphidium subarcticum* Cushman, 6-7. *Elphidium crispum* (Linné),
8. *Polystomelina discorbinoides* Yabe and Hanzawa, 9. *Elphidium jenseni* (Cushman),
10. *Criboelphidium yabei* (Asano), 11. *Criboelphidium oregonensis* (Cushman and Grant)

Plate 7

1, 8. *Astrononion hamadaense* Asano,
2. *Elphidium excavatum* (Terquem) forma *clavata* Cushman,
3. *Nonion japonicus* Asano, 4. *Melonis uchioi* Hasegawa, 5. *Pullenia apertura* Cushman,
6. *Melonis pompilioides* (Fichtel and Moll), 7. *Lenticulina nicobarensis* (Schwager),
9. *Nonionellina labradorica* (Dawson)

Plate 8

1. *Planoglabratella?* sp. C, 2. *Planoglabratella australensis* (Hellon-Allen and Earland),
3. *Planoglabratella opercularis* (d'Orbigny), 4. *Planoglabratella subopercularis* (Asano),
5. *Glabratella mirabiris panamensis* Seiglie and Bermúdez,
6. *Planoglabratella patelliformis* (Brady), 7-8. *Glabratella* sp. A,
9. *Glabratella pulvinata* (Brady), 10. *Glabratella* sp. B

Plate 9

1. *Planoglabratella subopercularis* (Asano), 2. *Epistominella pulchella* Husezima and Maruhasi,
3. *Ammonia beccarii* (Linné), 4. *Islandidilla norcrossi* (Cushman),
5. *Islandiella japonica* (Asano and Nakamura), 6. *Islandiella yabei* (Asano and Nakamura),
7. *Islandiella helanae* Feyling-Hanssen and Buzas,
8. *Islandiella sublimbata* (Asano and Nakamura),
9. *Pseudoparella takayanagii* (Iwasa), 10. *Pseudoparella naraensis* Kuwano

Plate 10

1. *Quinqueloculina* sp.G, 2. *Pateoris hauerinoides* (Rhunbler),
3. *Quinqueloculina costata* d'Orbigny, 4. *Quinqueloculina kuromatunaiensis* Asano,
5. *Quinqueloculina agglutinata* Cushman, 6. *Quinqueloculina* sp.F

Plate 11

1. *Cycloforia contrita* (d'Orbigny), 2. *Quinqueloculina akneriana* d'Orbigny,

3. *Quinqueloculina* sp. C, 4. *Cribrulinoides curta* (Cushman), 5. *Quinqueloculina* sp. B,
6. *Quinqueloculina yessoensis* Asano

Plate 12

1. *Quinqueloculina akneriana* d'Orbigny, 2. *Quinqueloculina sawanensis* Asano,
3. *Triloculina suttuensis* Asano, 4. *Triloculina tricarinata* d'Orbigny, 5-6. *Pyrgo ezo* Asano























